

Mexico's Biocultural Diversity in Peril

Omar Vidal^{1*} & Richard C. Brusca²

1. Bosque de Granados 141, Col. Bosques de las Lomas, Ciudad de México 11700, México; vidalpomar@gmail.com
2. Department of Ecology & Evolutionary Biology, University of Arizona, United States of America; rbrusca@desertmuseum.org

* Correspondence

Recibido 16-XII-2019. Corregido 26-II-2020. Aceptado 27-III-2020.

ABSTRACT. **Introduction:** Places with high species diversity have high linguistic diversity, whereas areas with low species diversity tend to have low linguistic diversity. **Objective:** To characterize the intriguing relationship between biological and cultural diversity, a correlation that has been discussed at a global scale, but here tested for the first time in Mexico. **Methods:** We compiled exhaustive databases on both endangered species and endangered languages, and reviewed available literature on Mexico's biocultural diversity with a focus on endangered and critically endangered species and languages. **Results:** With 364 living languages, Mexico is the world's fifth most linguistically diverse country, but 64 of these languages are facing a very high risk of disappearance and 13 have already disappeared. Mexico is also the fourth most biologically diverse country, but 1 213 species of its flora and fauna are threatened with extinction and at least 127 species were recently extinct. **Conclusions:** Indigenous peoples are custodians of much of the world's biocultural diversity. As the world grows less linguistically and culturally diverse, it is also becoming less biologically diverse. Mexico's biological and linguistic diversity show strong geographic overlap, with the states of Oaxaca, Chiapas, Veracruz, Guerrero, and Michoacán harboring most species and most languages. Similarly, Mexico's biodiversity hotspots mirror language hotspots, and areas with the highest number of endangered species overlap with areas where the endangerment of languages is also the highest.

Key words: languages, linguistics, traditional knowledge, indigenous people, biodiversity, Mexico, endangered species, extinction.

Vidal O., & Brusca, R.C. (2020). Mexico's Biocultural Diversity in Peril. *Revista de Biología Tropical*, 68(2), 669-691.

The United Nations Convention on Biological Diversity defines “indigenous” as those people who have historical continuity with pre-invasion and pre-colonial societies, that have developed on their own territories, and who consider themselves distinct from other sectors of society now prevailing in those territories. To raise awareness of their importance for humanity, the United Nations designated 2019 as the “International Year of Indigenous Languages.”

Biocultural diversity encompasses the link between biological diversity and humankind's cultural diversity, and identification of

ecological conservation hotspots and linguistics are the two pillars of biocultural diversity analyses. Trends in biological diversity and cultural/linguistic diversity parallel one another (Maffi, 2001, 2005) and there is a documented correlation between the two (Mühlhäusler, 1995; Harmon, 1996; Nettle & Romaine, 2000; Oviedo, Maffi, & Larsen, 2000; Moore et al., 2002; Sutherland, 2003; Stepp et al., 2004; Fincher & Thornhill, 2008; Gorenflo, Romaine, Mittermeier, & Walker-Painemilla, 2012). Research across both continental and regional scales have identified patterns of co-occurrence



of linguistic and biological diversity around the world, and nations/areas with high biodiversity also tend to have high linguistic and cultural diversity (Toledo, 1994; Harmon, 1995, 1996; Nettle, 1996; Nettle & Romaine, 2000; Stepp et al., 2004; Loh & Harmon, 2005; Toledo & Barrera-Bassols, 2008; Gorenflo et al., 2012). The countries with the highest linguistic diversity (jointly having 54 % of all living languages) are Papua New Guinea, Nigeria, India, Mexico, Cameroon, Australia, the Democratic Republic of the Congo, and China (Harmon, 1995); and the most bioculturally diverse countries as measured by their centers of biological, linguistic, and agricultural diversity, as well by the presence of indigenous peoples, include those plus Indonesia, Brazil and Peru (Toledo & Barrera-Bassols, 2008).

The conceptual framework of biocultural diversity draws upon a common interest in understanding and preserving the relationships between biological, linguistic, and cultural variety and range. For indigenous people, biological, cultural, and linguistic diversities are intrinsically linked, as are environment and development. Biocultural diversity is often used as an index, or measure, to assess geographical regions in terms of the status or linkages between biological, cultural and linguistic diversity (Harmon, 1996; Sutherland, 2003; Harmon & Loh, 2010; Gorenflo et al., 2012). Languages, like genes, are “documents of history,” and a vast amount of information about our past is inscribed in the content and structure of the approximately 7 100 languages that are spoken today (Gray, Quentin, & Greenhill, 2018). Present-day indigenous cultures and languages are an expression of the long historical legacy of interrelationships between humans and nature (Toledo & Barrera-Bassols, 2008). Conservation practitioners should also embrace biocultural approaches for social justice, legal, and practical reasons (Gavin et al., 2013, 2015). This view was convincingly articulated in the Indigenous Peoples International Declaration on Self-Determination and Sustainable Development at the 2012 United Nation’s Rio+20 Summit, which reaffirmed that cultural

belief systems and worldviews of indigenous people are fundamental to biodiversity protection and sustainable development.

Species are the basic units of biodiversity, while languages serve to measure the diversity of cultures, and the striking parallels in their evolution suggest that nature and culture evolved in similar fashion (Harmon, 1996). In practical terms, there is an emphasis on language over other aspects of culture and identity because it strongly circumscribes an indigenous group. *Homo sapiens* displays a remarkable linguistic diversity that correlates strongly to areas of plant and animal diversity—and both may be moderated by similar environmental factors, such as temperature and rainfall. Both domains also involve the transmission of discreet heritable units: genes in biology, and socially-transmitted units such as words and morphosyntax in linguistics. Both can be altered as they pass through generations and thus, they display a hierarchical relationship over time. Languages, like biological species, are therefore related through nested patterns of descent allowing their evolutionary history to be depicted in branching patterns, or trees (Dunn, 2014). Of course, some words do enter languages by way of diffusion, or “borrowing” from other cultural sources (Campbell, 1997), and these instances are comparable to horizontal gene transmission in the scheme of biodiversity assessment and phylogenetics. Nonetheless, just as the historical evolution of species can be inferred by phylogenetic analysis, so the history of linguistic diversity can be estimated by phylogenetics (e.g., Bouckaert et al., 2012; Birchall, Dunn, & Greenhill, 2016; Gray et al., 2018). In both cases, the phylogenies can be overlain by areas of endemism to produce dated phylogeographic reconstructions. In the case of cultures and linguistics, this amounts to what has been called “virtual archeology.”

Three types of language diversity have been recognized (Gavin et al., 2013). *Language richness* refers to the number of languages within a given area. *Phylogenetic language diversity* is the minimum total length of all branches needed to span a set of “language



taxa” on a phylogenetic tree. *Linguistic disparity* refers to the range of expression in a language trait within a clade. Languages differ on a multitude of structural levels, including phonology, morphology, and syntax. There is a strong latitudinal gradient in both language richness and biological richness (Gavin et al., 2013, 2015).

It has been proposed that human languages reached their maximum number (estimated at 12 000) at the end of the Pleistocene, directly predating the rise of agriculture (Harrison, 2007). In the non-sedentary hunter-gatherer societies of the time, the dominant force in language creation is likely to have been fissioning mechanisms (Hamilton, Milne, Walker, Burger, & Brown, 2007), although such historical “splits” might be blurred by inter-language diffusion, or borrowing, as noted above (e.g., Campbell, 1997). The number of languages has been in decline since the Neolithic as agricultural groups have spread, replacing hunter-gatherers, and population movements have tended to reduce global language diversity (Nettle, 1999a, 1999b). A substantial body of theoretical work suggests that the spread of politically complex agricultural societies was a dominant factor in the reduction of language diversity (Renfrew, 1994).

After examining nearly 7 000 languages spoken on Earth today against the planet’s biological diversity and biodiversity hotspots, Gorenflo et al. (2012) found that more than 4 800 languages occur in regions containing high biodiversity, and 3 202 languages occur in 35 previously defined biodiversity hotspots. Of the languages found in biodiversity hotspots, 1 553 are spoken by 10 000 or fewer people, and 544 are spoken by 1 000 or fewer people.

Greater geographic heterogeneity, as measured by more-diverse habitats or higher topographic complexity, is commonly correlated with greater species diversity (e.g., Kerr & Packer, 1997). A link between geographic heterogeneity and diversity also appears to exist for languages. Environmental variables have been noted as important predictors of linguistic diversity patterns, and historical processes of

co-evolution of small-scale human groups with their local ecosystems have been proposed (Harmon, 1996; Maffi, 1998).

Three narratives have been offered to explain the correlation between biological and cultural diversity (Maffi, 2001): *geographic determinism* (both species and languages diversify in heterogeneous landscapes with geographic barriers); *ecological determinism* (linguistic diversification occurs in response to high biodiversity as cultural groups encode their knowledge of rich biotas); and *historic determinism* (areas of high linguistic diversity today are “residual,” persisting because of their geographic isolation from extensive agricultural development, implying that now-impoverished areas were once more diverse).

Today, linguistic diversity and biological diversity face similar threats and are both in crisis because of human population growth and increasing consumption (Loh & Harmon, 2014). One of the biggest challenges for bio-cultural conservation is ensuring that indigenous peoples obtain recognition of their rights to the resources found on their lands and territories on which they depend on for their economic, spiritual, cultural, and physical well-being (Inter-American Commission on Human Rights, 2009), and that these translate into tangible policies and action at the national and international level. Which is, by no means, an easy task since it has been estimated that about 12-20 % of areas under human management worldwide are indigenous lands (Toledo, 2001a).

Different cultures perceive and value biodiversity in different ways because of their distinct heritage and experience. Posey (1999) argued that cultural diversity parallels ecological diversity, and local traditional adaptations are often the most environmentally sound. Through time, as local communities interact closely with their environment and modify it as they adapt to specific ecological niches, they acquire detailed knowledge of the environment and how to manage it for their long-term benefit. Languages thus carry deep eco-cultural knowledge, and embedded in indigenous



languages is knowledge about habitats, plant life, animal behavior, conservation methods, and many other aspects of the natural world (Harrison, 2007).

The world's inhabitants today speak around 7 100 languages, patterned unevenly across the Earth. But, roughly half of the world speaks only 24 languages, and have from tens to hundreds of millions of speakers, while the other half of the population speaks the remaining 7 073 languages, of which around half have fewer than 10 000 speakers (Lewis, Simons, & Fennig, 2016; Simons & Fennig, 2018). Since 1950, 230 languages have disappeared and at least 3 000 have become endangered, of which 577 are critically endangered (78 have 10–50 speakers, and 146 have fewer than 10 speakers) (United Nations Educational, Scientific and Cultural Organization [UNESCO], 2010). And, according to UNESCO, nearly 40 % of the world's population lack access to education in a language that they speak or understand. In the next 30 years, over half of the world's languages will likely go extinct or be spoken by only a few old people (Harrison, 2007).

While much effort has and is being spent to stem the loss of plant and animal species, the loss of languages is receiving far less attention and is actually being ignored in many parts of the world, particularly in developing countries, for a variety of reasons. As the world grows less linguistically and culturally diverse, it is also becoming less biologically diverse. Biologists estimate an annual loss of species at 1 000 times or greater than historic rates, and linguists predict that 50–90 % of the world's languages will disappear by the end of this century (Nettle & Romaine, 2000; Gorenflo et al., 2012).

More than 70 % of the world's biodiversity is found in only 17 countries, of which 15 are in the developing world and 6 are in Latin America (Mittermeier, Myers, Gil, & Mittermeier, 1999). The most species-rich, human-accessible environments are tropical rain forests and deciduous forests, coral reefs, and large tropical lakes (Millennium Ecosystem Assessment [MEA], 2005). Even though the world's tropical forests occupy only 7 %

of the land area, they have been estimated to contain over half the world's species (Oviedo et al., 2000; Corlett & Primack, 2010). Of the 93 579 species of plants and animals assessed by the Red List of the International Union for the Conservation of Nature (IUCN) until 2018, more than 26 000 (28 %) are threatened with extinction, including 41 % species of amphibians, 35 % of reptiles, 25 % of mammals, 13 % of birds, 7.5 % of bony fishes and 63 % of cycads. Those estimates, however, are very conservative given that the species evaluated only represent about 4 % of the nearly two million living species described to date (Brusca, Moore, & Shuster, 2016; Primack & Vidal, 2019), and far less than the estimated total number of species on Earth (estimates range from around 10 to 100 million species of prokaryotes, protists, plants, fungi, and animals) (Mora, Tittensor, Adl, Simpson, & Worm, 2011; Brusca et al., 2016). At least 479 species of vertebrates and 116 species of plants have gone extinct in modern times, due mainly to habitat destruction and fragmentation, over-exploitation, and invasive species. Although the global extent of protected land has roughly doubled in size since the 1992 Earth Summit in Rio de Janeiro, with more than 202 000 protected areas now covering 14.7 % of the world's terrestrial area, $6 \times 10^6 \text{ km}^2$ (32.8 %) of that protected land globally is under intense human pressure (Jones et al., 2018).

In this paper, we test the hypothesis that a pattern of co-occurrence of biological and cultural diversity exists in Mexico. We do this by examining the available data on species diversity and language diversity across all Mexican states, and by looking for areas of overlap between regions of high diversity and extinct/threatened/endangered diversity in species and languages.

MATERIALS AND METHODS

Much has been written on Mexico's biodiversity (for a review see Sarukhán et al., 2017). However, little has been published in the peer-reviewed literature on its rich linguistic



diversity. We extensively reviewed the available literature, including government reports, on Mexico's biocultural diversity with a particular focus on endangered and critically endangered species and languages to look for overlapping geographical distributions and threats. Most of the information we use regarding the languages of Mexico comes from an exhaustive appraisal of reports by Mexico's National Institute of Indigenous Languages (Instituto Nacional de Lenguas Indígenas [INALI], 2012) and official published government records (Diario Oficial de la Federación, 2008, 2010a). From these sources, we compile a robust set of data, though much of it is qualitative (not quantitative).

Our ranking criteria for biological species follow the IUCN, and we consider a species *critically endangered* when it faces an extremely high risk of becoming extinct, and *endangered* when it faces a very high risk of becoming extinct (IUCN, 2018). We follow two closely aligned criteria for defining endangered languages. The first, more detailed criterion, from INALI (2012), considers that a language has *very high risk* of disappearance when the total number of speakers is less than 100, the number of child speakers is less than 25 %, and the speakers are less than 30 % of the population in all localities; and a language is at *high risk* of disappearance when the total number of speakers is less than 1 000 but higher than 100, the number of child speakers is less than 25 %, and the number of speakers in just one locality is less than 30 %. The second criterion considers a language *critically endangered* when the number of speakers is less than 50, and *endangered* when the number of speakers is less than 250 (Loh & Harmon, 2014; Simons & Fennig, 2018). Both of these criteria are in line with the degrees of endangerment established by UNESCO (2010).

A linguistic variant has structural and lexical differences when compared with other variants in the same linguistic group. And those that speak a variant have a sociolinguistic identity that contrasts with the sociolinguistic identity of those that speak other variants within the same linguistic group (Diario Oficial de

la Federación, 2008). Those differences could be sounds, words, meanings, and uses that result in speakers of one variation being unable to understand speakers of other variations. Variants are therefore often seen as equivalent to languages (INALI, 2012) and are considered as such in this paper.

RESULTS

Biological Species

Mexico harbors 10-12 % of the world's biological diversity and is ranked as the fourth megadiverse country after Indonesia, Brazil and Colombia (Mittermeier & Mittermeier, 1992; Mittermeier et al., 1999; Toledo, Boege, & Carrera-Bassols, 2010; Sarukhán et al., 2017). So far, 118 030 (66.44 %) of the estimated 177 641 species of animals and vascular plants of Mexico have been described (Comisión Nacional para el Conocimiento y Uso de la Biodiversidad [CONABIO], 2008; Sarukhán et al., 2017): 23 314 species of vascular plants, 564 mammals, 1 150 birds, 908 reptiles, 399 amphibians, 2 763 fishes (2 224 marine), 11 472 non-arthropod invertebrates, 11 185 non-insect arthropods, and 66 275 insects. These estimates include terrestrial, freshwater, and coastal marine species.

As in other parts of the world, tropical regions in Mexico contain more species than temperate areas; although there are certain groups that are particularly diverse in arid zones, such as cacti, of which at least 677 species are found in the country, 518 of them being endemic (Dávila et al., 2002; Villaseñor, 2016). The states of Oaxaca, Chiapas, Veracruz and Guerrero (all in the South/Southeast and close to Guatemala and Belize) and Michoacán are home to the bulk of the country's biodiversity (see also Toledo et al., 2001). Mammal, reptile and amphibian species richness is much higher in Oaxaca, Chiapas, Veracruz and Tabasco (CONABIO, 2008) and although it varies among different groups, tropical humid regions appear to have fewer endemic species



than arid and semiarid regions (CONABIO, 2016). Oaxaca alone is home to almost half of Mexico's known species of vertebrates, 19 % of the known invertebrates, and 40 % of the known plants (Gonzalez-Perez, Briones-Salas, & Alfaro, 2004).

At least 127 species of Mexican flora and fauna (58 % of them endemic) are known to have gone extinct through 2008 (Table 1). However, since most recorded extinctions have occurred on islands and in continental lagoons and rivers, and include only flowering plants and vertebrates, the actual number of extinctions (including other habitats and taxa) is probably much higher. There are today 1 213 Mexican species (many endemic) threatened with extinction (Diario Oficial de la Federación, 2010b; IUCN, 2018; **Supplemental Material**): 94 (73 endemic) species of mammals, 66 (24) birds, 97 reptiles, 219 (182) amphibians, 181 fishes, 8 mollusks, 98 other invertebrates, and 450 plants (including 12 conifers, 41 cycads, 133 cacti and 22 magnolias). Unsurprisingly –given the extent of habitat destruction and fragmentation in Oaxaca, Chiapas, Veracruz, Guerrero, Michoacán, and states adjacent to the Gulf of California, including its many islands and islets— threatened hotspots are located mostly in those regions.

Languages

Today, 25.7 million (21.5 %) of Mexico's inhabitants are indigenous peoples, and 7.4 million (6.5 % of all Mexicans older than three

years) speak an indigenous language (Instituto Nacional de Estadística y Geografía [INEGI], 2010, 2015); although in some states such as Chiapas, Oaxaca and Yucatán that percentage is nearly 30 % (CONABIO, 2016). Indigenous land represents approximately 14.3 % (or 28 million hectares) of the country's total territory (Boege, 2008, 2009) and nearly half of the most important watershed headwaters are occupied by indigenous peoples. About a third of Mexico's protected natural areas (at the federal level) have indigenous populations living within them, and approximately 70 % of Mexico's indigenous land is under some sort of priority for the conservation of its rich biological resources (Toledo et al., 2010). There are 64 172 named localities with indigenous people and a quarter of Mexico's social properties are located within 4 786 *ejidos* (communal land, which in Mexico is land expropriated from owners of large tracts and redistributed for shared use as farmland, especially to poor populations, in accordance with the Agrarian Reform Act of 1917) and 1 258 agrarian communities in territories own by indigenous people (INEGI, 2015). The states of Oaxaca, Chiapas, Veracruz, Guerrero, Hidalgo, Estado de México, and Yucatán are home to 77 % of the Mexico's indigenous people, while the states of Coahuila, Colima and Zacatecas have the lowest indigenous populations.

Today, with 364 living languages, Mexico is the world's fifth most linguistically diverse country after Papua New Guinea, Indonesia,

TABLE 1
Mexico's recorded extinct species of fauna and flora (CONABIO, 2008; Sarukhán et al., 2017)

Taxa	Extinct Species	Endemic Species	States/Islands
Plants	26	5	Hidalgo, Veracruz, Jalisco, Isla Guadalupe
Fish	38	20	Nuevo León, Coahuila, Jalisco, Durango, México, Veracruz
Amphibians	29	29	Oaxaca, Veracruz, Guerrero, Querétaro, Hidalgo, Durango
Birds	19	11	México, Sonora, Coahuila, Michoacán, Colima (Isla Benedicto), Baja California (Islas Todos los Santos), Isla Guadalupe, Isla Socorro
Mammals	15	9	Baja California (Islas Todos los Santos, Coronado, Turner, and San José), Nayarit (Archipiélago las Marias), Sonora (Isla San Pedro), Isla Ángel de la Guarda, Isla San Juanito
Totals	127	74	



Nigeria and India (*Diario Oficial de la Federación*, 2008; Toledo et al., 2010; INALI, 2012). Mexico's indigenous languages belong to 11 linguistic families and 68 linguistic groups (*Diario Oficial de la Federación*, 2008; INEGI, 2010; INALI, 2012). This cultural richness is, however, threatened.

Four linguistic groups comprise the largest numbers of speakers in Mexico –Náhuas (1 376 000 speakers), Maya (759 000), Mixteco and Zapoteco (> 400 000)– while 22 groups each have less than 1 000 speakers. Oaxaca, Puebla, Chiapas, Veracruz and Guerrero are the states with the highest linguistic diversity; densities of languages in Oaxaca, Veracruz and Guerrero are comparable to areas in Papua New Guinea, the Himalayas, Nigeria, and Cameroon (CONABIO, 2008). Ninety percent of the people who speak the Cochimi-Yuman family (Cucapá, Paipai, Kumiai, Ku'ahl and Kiliwa) live in Baja California and Baja California Sur, all of which are at high or very high risk of disappearance. Cochimí is not a “modern” Yuman language, but one that likely split off from a Proto-Yuman tongue long ago (Mixco, 1978).

The status of speakers among ethnic groups differs greatly throughout Mexico. There are groups in which only the elders speak the language (in Mexico), as is the case of the Ixcatéco, Ayapaneco, Kiliwa, Paipái, Cucapá and Ku'ahl; and those for which adults speak the language, but children do not, such as Tlahuica, Mocho', Tuzanteco, Teko, Awakateko, Oluteco, Ayapaneco, Texistepequeño, Chocholteco, Kaqchikel, Ixil, and Ixil chajuleño (INALI, 2012). However, Kaqchikel and Ixil are languages probably represented today in Mexico by relatively recent immigration from Guatemala, where they remain strongly represented. At least 13 languages have gone extinct in Mexico: Pericú, Solteco, Naolan, Opata, Pochuteco, Cuitlatec, Pame (Southern), Tepecano, Tubar, Chiapaneco, Eudeve, Pochuteco and Cochimí. Of the 364 languages spoken today in Mexico, 64 are in very high risk of disappearance (less than 100 speakers survive) and 43 are in high risk of disappearance (less than 1 000 but more than 100 speakers survive (INALI, 2012) (Table 2, Fig. 1). Nine languages in Mexico are also considered by Simons and Fennig (2018) to be critically endangered (less than



Fig. 1. Mexico's languages at very high risk of disappearance shown in red (from INALI, 2012; coloring denotes areas in which the language is spoken), and its most biologically diverse states bordered in green.



TABLE 2
Extinct, critically endangered, and endangered languages in Mexico

EXTINCT				
Language (family)	Number of Speakers	State	Locality (municipality: villages) (1)	
Pericú (2)		Baja California	Los Cabos region	
Solteco (Oto-mangue) (3)	None	Oaxaca	Balsas River	
Cuitlatec (4)	None	Guerrero	San Juan Náolan, near Tula	
Náelan (unclassified) (5)	None	Tamaulipas	Pochutla	
Pochuteco (6)	None	Oaxaca	Arivechi, Bacanora, Nacorí, Onavas, Sahuaripa, Suaqui. Tecoripa is the traditional area	
Opata (Uto-azteca) (7)	None	Sonora	Jilíapan	
Pame, sureño (Oto-mangue)	None	Estado de México	(possibly Hidalgo)	
Tepecano (Uto-azteca) (8)	None	Jalisco	Villa Guerrero: Azqueltán (earlier Atzqueltán), San Martín de Bolaños on Río Bolaños	
Tubar (Uto-azteca) (9)	None	Chihuahua	near the border with Sonora and Sinaloa	
Chiapaneco (Oto-mangue)	None (10)	Chiapas	El Bosque, Las Margaritas, Ocosingo, Atenque and Sabanilla municipalities	
Eudeve (11)	None	Sonora (12)		
Pochuteco (Uto-azteca) (13)	None	Oaxaca	Pochutla	
Cochimí (Cochimí-yumana) (14)	None (15)	Baja California	Noth of Loreto	
CRITICALLY ENDANGERED (less than 50 speakers and no children -5 to 14 years of age- speak the language)				
Kiliwa (Cochimí-yumana)	2 (16)*	Baja California		
Awakateco (Maya)	3*	Campeche	Ensenada: Arroyo de León (Ejido Kiliwas), Ejido San Francisco R. Serrano (Valle San Matías), Ensenada, Francisco Zarco (Guadalupe), Juntas Neji, La Zorra, Lazaro Cárdenas (Valle de la Trinidad), Licenciado Gustavo Díaz Ordaz, Parcela Número 19 (Familia Castro Ejido Nata), Rancho las Pinzas	
Tuzanteco (Maya)	5*	Chiapas	Champotón: Maya Tecún I, Maya Tecún II, Santo Domingo Kesté	
Ayapaneco (Mixe-zoque)	8 (17)*	Tabasco	Tuzantán: Estación Tuzantán Jalapa de Méndez: Ayapa, El Carmen (La Ensenada), Vicente Guerrero Primera Sección	
Ixil neabajeno (Maya)	12*	Quintana Roo	Othón P. Blanco: Maya Balam	



Language (family)	Number of Speakers	State	Locality (municipality·villages) (1)
Zapoteco de Mixtepec (Oto-mangue)	14 (18)*	Oaxaca	San Cristóbal Amatlán: San Agustín Mixtepec
Ku'ál (Cochimí-yumana)	20 (19)*	Baja California	Misión de Santa Catarina, Comunidad Indígena de Santa Catarina, El Aguaajito (Mat Chip) [Rancho Matchip], Rancho Escondido, Rancho Wikualpuk (El Ranchito), Ensenada
Ixcateco (Oto-mangue) (20)	21 (21)*	Oaxaca	Oaxaca de Juárez, Santa María Ixcatlán
Kaqchikel (Maya)	35*	Campache	Campache: Quetzal-Edzná I, Quetzal-Edzná II, Los Laureles. Champotón: Maya Tecún II, Santo Domingo Kesté
Zapoteco de San Felipe Tejalapám (Oto-mangue)	50 (22)*	Quintana Roo Oaxaca	Quintana Roo: Othón P. Blanco: Kuchumatán, Maya Balam, San Isidro la Laguna San Felipe Tejalapám: Arroyo Central, Cañada Cera, Conejo (Rancho El Conejo), El Chilar, El Copalito, El Cuchanal, El Paredón, El Roble, La Aurora, La Unión, Llano Ceniza, Llano El Coyote, Los Paderones, San Felipe Tejalapám
ENDANGERED (less than 250 speakers no children -5 to 14 years of age- speak the language)			
Ixil chajuleño (Maya)	52	Campache	Campache: Los Laureles, Quetzal-Edzná
Zapoteco de Asunción Tlacolula (Oto-mangue)	53	Quintana Roo Oaxaca	Quintana Roo: Othón P. Blanco: Kuchumatán Asunción Tlacolula: Asunción Tlacolula, Primera Sección Norte
Obliteco (Mixe-zoque)	60*	Veracruz	Oluta: Correa de Abajo, El Chorro, El Mirador, El Ángel Gabriel, Los Laureles, Oluta, Tenejapa
Chocholteco del oeste (Oto-mangue)	62	Oaxaca	Ocotlán: Boquerón, San Antonio Nduyaco, Tierra Colorada, Unión Palo Solo. San Francisco Teopan: Concepción Buenavista, Santiago Ihuatlán Plumas, Tepelmeme Villa de Morelos, Tacotepec Plumas.
K'iche' (occidental) (Maya)	65	Campache	San Miguel Tulancingo: Agua Dulce, Buena Vista, El Coatillo, El Españo, Gasuchío, Loma Larga, Rancho Marino Sánchez, San Miguel Tulancingo, San Pedro Nopalá: San Mateo Tlapilepec, Santa María Jicotlán, Santa Magdalena Jicotlán: San Mateo Tlapiltepec, Santiago Tepetlapa. Teotongo: El Progreso, El Tecomate, Guadalupe, La Luz
K'iche' (central) (Maya)	65	Campache	Champotón: Santo Domingo Kesté
Teko (Maya)	71*	Chiapas	Amatenango de la Frontera: Amatenango de la Frontera, Barrio Nuevo, Chiquisbil, El Porvenir, Granadilla, Las Marías, Nuevo Amatenango, Sabinalito. Frontera Comalapa: Nuevo Mazapa.
			Mazapa de Madero: Bacanton Altamirano Uno, Chimalapa, Mazapa de Madero, Nuevo Paraíso, Reforma, Tierra Blanca, Valle Obregón, Veracruz, Villa Hidalgo.



Language (family)	Number of Speakers	State	Locality (municipality· villages) (1)
Zoque del sur (Mixe-zoque)	84	Chiapas	Ocozocuautla: El Edén, El Pedregal, Nuevo Chapultenango, Salina Cruz, San Antonio, San Luis las Maravillas, San Pedro, Velasco Suárez Uno. Tuxtla Gutiérrez: Copoya, Julio César Ruiz Ferro Segunda Sección, Lindavista, Los Crisantemos.
Pápago (Pima alto) (Yuto-nahua)	94	Sonora	Altar, El Cubabi, El Cumariito. Caborca: El Sofí, Heroica Caborca, Juárez, La Escondida, Las Norias (De Romero), Puerto Lobos, San Francisco Papagos (San Francisquito), San Pedro, Santa Eduwiges (La Cachora), SPR Mayobampo (Rancho Corralero).
K'iche' (oriental) (Maya)	96	Campêche	General Plutarco Elías Calles: Ejido Luis Echeverría Álvarez, El Chamizal, El Desierto de Sonora, Quijovaec, San Antonio, Santa Rosa (La Angostura), Sonoya. Campeche: La Libertad. Champotón: Maya Tecum II, Santo Domingo Kesté.
Kuapá (Cucapá) (Cochimi-yumana)	119 (24)	Chiapas	Chiapas. Las Margaritas: San Mateo Zapotal. Marqués de Comillas: Boca de Chapul, Pico de Oro.
Kickapoo (Álgica) (23)	105	Quintana Roo	Quintana Roo. Othón P. Blanco, Kuchumatán, Maya Balam, San Isidro la Laguna.
Mexicano de occidente (Yuto-nahua)	107	Coahuila	Múzquiz: Ciudad Melchor Múzquiz, El Nacimiento de los Kikapíes (Nacimiento de la Tribu Kikapoo)
		Jalisco	Cuauitlán de García Barragán: Ayotlán, Cortácpo, Cuauitlán de García Barragán, Cuzalapa, Chacala, Chanquehuil, El Chico, La Guaca, La Rosa, La Rosita, Los Encinos, Los Sauces, Los Terreros, Mojoneras, Paso Real, Plan de Méndez, Plan de San Antonio, Rancho Viejo, San Miguel, Santa Rosa, Tierras Blancas, Tierras Negras
		Baja California	Baja California: El Mayor Cucapá, Pozas de Arvizu. Mexicali: Campo Camerina (Colonia Terenos Indios), Campo del Prado (Colonia el Mayor), Campo Flores, Campo Sonora (Colonia Terenos Indios), Colonia la Puerta, Comunidad Indígena Cucapá el Mayor [Ejido el Mayor], Ejido Cucapá Mestizo, Ejido Doctor Alberto Mota (El Indiviso), Ejido Durango, Ejido México, Familia Regalado (Ejido Sonora 2 Campos Nuevos), La Casa de las Curvas (Colonia el Mayor), Mexicali, Sainz Domínguez (Colonia el Mayor), San Felipe.
Mexicano alto de occidente (Yuto-nahua)	127	Sonora	Sonora. San Luis Río Colorado: Pozas de Arvizu (La Reserva).
Mocho' (Maya)	141*	Nayarit	Acaponeta: Acaponeta, Agua Tendida, Buenavista (Las Paredes), El Duraznillo, El Narrajo, El Resbalón, Huancaxtle, La Guásima, La Laguna (La Lagunita), La Paloma Nueva Reforma, Mesa las Arpas, Paredes, Rancho los López, Rancho de Ernesto Aguilar, San Diego de Alcalá, San Diego el Naranjo, San Dieguito de Abajo, San José de Gracia, Santa Cruz, Sayocota, Sayullilla, Unidad Habitacional, Zácatecas. Del Nayar [El Nayar]: Dolores, Los Chapiles, Santa Teresa, Tierras Cuevas Motozintla: Motozintla de Mendoza



Language (family)	Number of Speakers	State	Locality (municipality: villages) (1)
Pima del norte (Yuto-nahua)	155	Chihuahua	Madera: Agua Amarilla, Ciudad Madera, Ejido el Lago, El Cable, El Campo Seis, El Cordón, El Cuatro, El Largo, El Pedregal, El Potrero, El Presón, El Represito, El Río Chiquito, El Táscate, El Yerbanís, Junta de los Arroyos [Junta de los Ríos], La Bolada, La Ciénega, La Nopalera, Las Espuelas, Las Lajitas, Los Arbolitos, Madera, Mesa Blanca, Mesa el Tecolote, Mesa de La Simona, Mesa del Garabato, Rancho Huapoca, San Isidro, San Juan de Enmedio. Moris: Bermúdez, Los Cien Pinos
Pima del sur (Yuto-nahua)	159	Chihuahua	Ocampo: Basaseachi Pinos Altos, Santo Domingo, Sapareachi, Tierritas. Temosachí: Agua Caliente, Casa Blanca, Ciénega Blanca, Cordon de Enmedio (Paraje Piedra Colorada), El Arbolito, El Llorón, El Nogal, El Saucito, El Terrero (Piedras Azules), Janos, La Ciénega, La Ciénega Blanca [Ciénega Blanca], La Guajolota, La Omela, La Providencia, La Salitrera, La Vinata, Las Tierritas, Los Hornitos, Nabogame, Peñasco Blanco, Piedra Blanca, Piedras Azules, San Antonio, San Ignacio, Temósachi, Tierras de María, Yahuirachi, Yepachic.
Paipai (Cochimi-yumana)	162 (25)	Baja California (26)	Ensenada: Arroyo de León (Ejido Kiliwas), Camanal, Cañón de la Parra, Comunidad Indígena de Santa Catarina, Ejido 18 de Marzo (El Alamo), El Agnajito (El Mat Chip), El Alamar, El Pinacate [Chiknan], El Ranchito [Wikwalipuk], El Sauzal, Ensenada, Ex Hacienda Sinaloa, Héroes de la Independencia (Llano Colorado), Lázaro Cárdenas (Valle de Trinidad), La Huerta, La Vinata [Wipuk], Leyes de Reforma (El Rodeo), Misión Santo Domingo, Ojo de Agua Colorada, Poblado Héroes de Chapultepec, Poblado Puerta Trampa, Pórticos del Mar, Rancho Xonuko, Rancho Agua de Vida, Rancho el Sauco, Rancho Escondido, Rancho las Canoas, Rancho Mariseal, Rancho San Belén, Rancho Santa Martha, Real del Castillo Nuevo (Ojos Negros), Rinconcón de Santa Catarina, San Isidoro, Sauce Largo [Yokagulg], Sauce Solo [Yokazis], Úrsulo Galván, Valle de la Trinidad, Yokakul
Zapoteco de la Sierra sur, noroeste bajo (Oto-mangue)	164*	Oaxaca	Villa Sola de Vega: Agua del Sabino, Arroyo de la Rosa (El Aguacate). Barrio de Santa Anita (Sección Tercera), El Anis (San José El Anis), El Duraznillo, El Mangalito, El Nido, El Progreso, Guayabo, La Ciénega de la Palma, La Labor, La Nopalera, La Siempre Viva, La Singuera, Las Juntas (El Puente), Las Trancas, Llano Mazorca, Llano Monte, Llano de León, Loma de Ocoté, Rio Espejo, Rio Húmedo, Río Mantecca, Río de Boca, San Felipe Zapotitlán, San Juan Viejo, San Sebastián de las Grutas, Villa Sola de Vega
Chocholteco del este (Oto-mangue)	180	Oaxaca	San Juan Bautista Coixtlahuaca: El Capulin (Sección Primera), El Tepozon (Sección Segunda), El Zapotal (Sección Tercera), La Mulata, Santa Catarina Ocotlán. San Miguel Chicahua: Llano Seco.
Chocholteco del sur (Oto-mangue)	200	Oaxaca	San Miguel Huautla: Ocotlán. Santa María Nativitas: Barrio Nicolás, Barrio Santiago, El Mirador, El Porvenir, Loma del Tepejillo, Pie del Cordoncillo, Primera Sección (Santa Cruz), San José Monte Verde, San Pedro Buenavista, Santa María Nativitas.



Language (family)	Number of Speakers	State	Locality (municipality: villages) (1)
Zapoteco de Valles, oeste (Oto-mangue)	211	Oaxaca	Magdalena Mixtepec: Magdalena Mixtepec, Santa Catalina Mixtepec. San Bernardo Mixtepec: Asunción Mixtepec, Barrio del Carmen, Barrio del Rosario, La Loma Grande, Rancho Teja, San Bernardo Mixtepec, San Jerónimo Tlaltetlán. San Pablo Cuatro Venados: El Espronjado, El Jabali, El Moral, El Polo, El Sacatón, La Nevería, Llano Manteica, Llano de Toro, Piedra Blanca, Río Jalapillas, Río Minas, San Cristóbal, San Pablo Chiatro Venados, Santiago Clavellina. San Pablo Huixtepec: San Pablo Huixtepec.
Mixteco del noroeste bajo (Oto-mangue)	220*	Oaxaca	San Bartolo Soyaltepec: Barrio del Calvario, Cerro Totocani, El Caballo Blanco, El Águila, Guadalupe Gavillera, Jataíno, La Unión Reforma, Río Verde, San Bartolo Soyaltepec, San Isidro Tejocatl, Santa Cruz Pipila. Santa María Chachopápm: San Agustín Monte Lobos, Santa María Chachopápm. Santo Domingo Tonaltepec: Cerro de Sol, Deque Inu (Cabeza de Cigarrero), Dequetoto, Río Verde, San Francisco Río Blanco, Santo Domingo Tonaltepec, Vista Hermosa, Yuquini. Santo Domingo Yanhuitlán: Santo Domingo Yanhuitlán, Tooxi, Yudayo.
Kumiai (Cochimi-yumana)	221 (27)	Baja California (28)	Ensenda: El Porvenir (Guadalupe), Ensenada, Francisco Zarco (Guadalupe), La Huerta, La Misión, Rancho Dinuwa, Rancho Plazaola, Real del Castillo Nuevo (Ojos Negros), San Antonio Necua (Cahada de los Encinos), San Salvador de Gálvez. Playas del Rosario: Comunidad Indígena San José de la Zorra (San José de la Zorra) [San José la Zorra], La Zorra, Primo Tapia. Tecate: Colonia Luis Echeverría Álvarez (El Hongo), Ejido Guadalajara 2 (El Gato), Ejido Nueva Colonia Hindú, El Álamo, El Testarazo, El Venado, Encino Solo, Escuela Mescuch [Mescuch], Familia López Gaspar, Hacienda Santa Verónica, La Ciénega, Rancho Cañada Verde, Rancho Jacuín, Rancho las Auras, Rancho las Priscillas, Rancho Limón, Rancho Neji Sección C [Juntas de Neji], Rancho Puerto el Roble, Santa Elena, Tecate, Valle de las Palmas.
Ch'ontal de Oaxaca bajo (Ch'ontal de Oaxaca)	223	Oaxaca	Santa María Ecatepec: La Reforma, San Juan Acaltepec, Santo Domingo Chontecomatlán, Santo Tomás Tejpan
Mixteco de San Miguel Piedras (Oto-mangue)	243*	Oaxaca	San Miguel Piedras: Chidoco (Chidoco de Juárez), Colorado, El Fresno, El Potrerito, Guadalupe Victoria, Río Minas (Río Minas Piedras), San Miguel Piedras
Zapoteco de San Bartolo Yutepet (Oto-mangue)	246	Oaxaca	San Bartolo Yutepet: Joaquín López Mariano (Rancho), Puerto San Bartolo, San Bartolo Yutepet mangue

Localities are from Diario Oficial de Federación (2008), status and notes are mainly from INALI (2012), Hammarström et al. (2018) and Simons & Fennig (2018). Critically endangered (< 50 speakers survive) and endangered (< 250 speakers survive) languages as in Simons and Fennig (2018); those denoted by * are in very high risk of disappearance (< 100 speakers survive) and high risk of disappearance (< 1 000 but > 100 speakers survive) following INALI (2012).



1. Localities included are those with historic settlements.
2. Also known as edúes y coras.
3. Related to Chatino and Zapoteco languages.
4. Last speaker died in the 1960s.
5. Extinct ca. 1950.
6. Boas (1917).
7. Last speakers reportedly died around 1930, but the 1990 census lists 12 speakers.
8. Formerly spoken by a small group of people. The last speaker was born in 1895 (Holt 2001) and no remaining speakers are known after 1972.
9. As many as 100 in 1970; see also Lionnet (1978).
10. Ethnic population is 32.
11. Also known as heve and dohema; extinct since about 1940; see also Lionnet (1986).
12. In her essay “El noroeste: Sonora” (<http://www.fundacionunam.org.mx/humanidades/eudeves-una-etnia-extinta/>), historian Isabel Verdugo de Juárez locates the territory occupied by Eudeves as: “In the north they were found along the mid-portion of the San Miguel River (Horcasitas River) in Saracachi, Cucurpe o Opodepe. In the south they we found at the springs along the Matepe River, the banks of Moctezuma River and part of Nevoma or Yaqui Rivers. This area encompassed the historical sites of Batuco, Tepupa, Bacanora, Soyopa and Tónachi.”
13. Extinct in 1954 (Boas 1917; Knab 1980).
14. From the first contact with Europeans 300 years ago, the Cochimíes have occupied the central part of the Baja California peninsula and the state of Baja California Sur. Although there was some minor dialect variation over the vast area where Cochimí was spoken, they were mutually intelligible (T. Bowen, pers. comm.). Originally, there were no large settlements and they were basically nomads. The guamas or sorcerers had an important position in the original culture; there were no writing or formal artistic expressions. They were gatherers and fishers, who did not practice agriculture nor have livestock.
15. Ethnic population is 150; this may include Kumiai in La Huerta who call themselves Cochimí; old Cochimí is extinct (Simons & Fennig 2018).
16. A total of 15 Kiliwa people survived in 2018 and the number of speakers is given as 29 (INALI 2012); however, in February 2018 one of the last three speakers died in Ejido Tribu Kiliwas, Valle de la Trinidad, near Ensenada.
17. Ethnic population of 16 in 2016.
18. It has been reported that in 2003 there were only two native speakers (the youngest then 72 years old) and three semi-speakers; the community apparently agreed to stop speaking Zapotec in 1965 (Beam de Azcona, 2004).
19. A total of 166 elder speakers was also reported (INALI 2010).
20. Reported as Xuani-Ixcateca by Molina Cruz (2010), who described efforts to save this language using video (see also <https://www.youtube.com/watch?v=IN2M-Rb7LM0>).
21. A total of population of nine and only a few elder speakers (in 2009) were reported (Simons & Fennig, 2018).
22. A total of 30 speakers, all over the age of 70, were reported in 1998 (Simons & Fennig, 2018).
23. The Kickapoo arrived in Mexico in the nineteenth century when, after the Anglo-Saxon invasion of their territory, they asked the Mexican government for a place to live; in exchange the government asked them to defend Mexican residents against the frequent attacks from the Comanche. Since then, the Kickapoo of Mexico have been known in the United States as the “Texas gang.”
24. Caccavari Garza (2014) reported less than 30.
25. Caccavari Garza (2014) reported less than 50.
26. According to Caccavari Garza (2014) most people are in Comunidad Indígena de Santa Catarina, but they are also present in San Isidoro, Valle de la Trinidad and Ejido Héroes de la Independencia.
27. Caccavari Garza (2014) reported less than 50.
28. According to Caccavari Garza (2014) mainly located in two municipalities and the communities of Juntas de Nejí en Tecate, San José de la Zorra, San Antonio Necua y La Huerta, Ensenada.

50 speakers survive), and 29 endangered (less than 250 speakers survive). The ones with the highest risk of going extinct in the immediate future are: Kiliwa (2 speakers survive), Awakateco (3 speakers), Tuzanteco (5 speakers), Ayapaneco (8 speakers), Ixil nebañeño (12 speakers), Zapoteco de Mixtepec (14 speakers),

Ku’al (20 speakers), Ixcateco (21 speakers), Kaqchikel (35 speakers), Zapoteco de San Felipe Tejalapám (50 speakers), Ixil chajuleño (52 speakers), and Zapoteco de Asunción Tlacolulita (53 speakers). Mexico’s 56 extinct and endangered languages occur in 16 states, most (77 %) in Oaxaca (16), Baja California



(7), Chiapas (6), Campeche (6), Sonora (4) and Quintana Roo (4) (Table 2, Table 3).

As a stark reminder, one language and one species, both located in Baja California, will probably go extinct within the next very few years. With only two speakers alive in late 2018, the Kiliwa is Mexico's most endangered language and is condemned to extinction (INALI, 2012; Caccavari Garza, 2014). And, with a population of fewer than two dozen individuals in 2018 (Comité International para la Recuperación de la Vaquita [CIRVA], 2019), probably even fewer today, the vaquita porpoise (*Phocoena sinus*), Mexico's only endemic and the world's most endangered marine mammal, is being driven to extinction by illegal fishing for the endangered and endemic totoaba (*Totoaba macdonaldi*) (Vidal, 1995; Brusca, Álvarez-Borrego, Hastings, & Findley, 2017; Thomas et al., 2017; Rojas-Bracho et al., 2018).

TABLE 3
States in Mexico with extinct or endangered languages

State	Number of extinct and endangered languages
Oaxaca	16
Baja California	7
Chiapas	6
Campeche	6
Sonora	4
Quintana Roo	4
Chihuahua	3
Jalisco	2
Tabasco	1
Veracruz	1
Guerrero	1
Tamaulipas	1
Baja California Sur	1
Mexico State	1
Coahuila	1
Nayarit	1

DISCUSSION

Places with high species diversity, especially tropical forests, tend to have high linguistic diversity, and areas with low species diversity, such as tundra and deserts, have low linguistic

diversity (Sutherland, 2003; Loh & Harmon, 2005). For instance, Papua New Guinea covers less than 1 % of the world's land area but is home to the world's third largest tropical rainforest (after the Amazon and the Congo) (Bryan, Shearman, Ash, & Kirkpatrick, 2010), and is also home to 6-8 % of the Earth's animal and plant species (two thirds of them endemic) and 12 % of the world's living languages (Papua New Guinea's Fifth National Report to the Convention on Biological Diversity, 2014; Simons & Fennig, 2018).

Indigenous people are custodians and landowners of much of the biodiversity worldwide. The world's 370 million indigenous people make up less than 5 % of the total human population, yet they manage or hold tenure over 25 % of the world's land surface that supports about 80 % of the global biodiversity (Raygorodetsky, 2018).

There are increasing numbers of examples of non-industrial people living in harmony with their natural environment, such as rural communities in Hawaii (McGregor, 1999); the Kayapo Indians of Middle Xingu Valley in Brazil (Posey, 1999); the Dai, an indigenous ethnic group in Southwest China (Shenghi, 1999); and the Sápara of Ecuador (Raygorodetsky, 2018). Numerous studies have proven how traditional ecological knowledge and practices have effectively served to protect and maintain natural environments (e.g., Posey, 1999; Berkes, 2001; Cunningham, 2001; Wiersum, 2004; Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services [IPBES], 2018). In addition, biocultural conservation can help secure the rights of indigenous and local people and help maintain a focus on social justice.

The current rate of extinction of languages worldwide, and loss of knowledge they contain, has no parallel in human history (Harrison, 2007). Language loss in some areas, such as the Americas, has reached 60 % over the last 35 years (Harmon & Loh, 2010). Indeed, the most rapid losses in linguistic diversity have occurred in this region, where 60 % of languages are threatened or have gone extinct since



1970. And this is not limited to developing countries. Policies in the United States, particularly between the 1870s and 1930s, greatly suppressed Native American languages and culture. It was only after years of activism by indigenous leaders that the U.S. Native American Languages Act was passed in 1990, which allowed for the preservation and protection of indigenous languages.

Languages can go extinct either because the entire population of speakers dies out, or because the speakers shift to a different language and forget their mother tongue due to the loss of intergenerational transmission. Much of the decline in linguistic diversity is a result of language shift away from small indigenous languages toward national or regional languages (Loh & Harmon, 2014). Along with the disappearance of languages, most of the rich traditional knowledge of these indigenous cultures is largely lost forever, including knowledge of the natural history of the world in which the language resided.

Languages are a critical measure of the cultural diversity of nations, while species richness is a measure of their biological diversity. Mexico offers an important case on the challenges to conserve both. With at least 118 species of economically important plants totally or partially domesticated by indigenous pre-Hispanic farmers, Mexico has been a global center of plant domestication. More than 15 % of edible vegetable species consumed in the world originated in Mexico (CONABIO, 2008; Perales & Aguirre, 2008; Bellón et al., 2009; Sarukhán et al., 2017). In addition, between 3 000-4 000 medicinal plant species are regularly used by Mexicans, and indigenous people in Mexico use 5 000-7 000 plant species in various cultural activities (Boege, 2008). It has been suggested that almost every species of plant and animal, type of soil, landscape, and mountain in Mexico has its match in a linguistic expression, knowledge category, practical use, mythical or religious meaning, or a known individual or collective experience (Toledo et al., 2001a, 2001b).

Mexico's early cultures relied heavily on natural resources, including the harvesting of coastal and estuarine resources in the Holocene. In the Baja California-Gulf of California region this included fishing and intensive hunting of dolphins and sea turtles beginning at least 10 000 years ago (Bowen, 1976, 2000, 2006, 2009; Felger & Moser, 1987; Felger, Nichols, & Seminoff, 2005; Fujita, 2006; Marlett, 2014). By around the year AD 1000, four important centers of indigenous socio-economic and cultural concentration surged in this region, which reflected changing patterns of harvesting of marine resources. First, during the early 16th and 17th centuries, the Pericú, who lived along the deserts of Los Cabos region in the South of the peninsula, were hunters and shell gatherers (*concheros*), and disappeared (together with their language, its phylogenetic classification remaining unknown) in the second half of 18th century (Fujita, 2006). The exploitation of marine mammals, the associated stone tools, and the construction of enormous shell mounds by the Pericú display striking similarities to the Chumash people of the Channel Islands in Southern California in the United States (Beer, Gonzalez, Huddart, Rosales-Lopez, & Lamb, 2008). Second, the Guaicuras and the Cochimís, the latter inhabiting the center and North of the Baja California peninsula, were also shell gatherers and fishers. Third, the Seris on the ("mainland") coastal Sonoran Desert were (and still are) fishers and gatherers. And fourth, the Yumas, Pimas Altos and Papagos in North-Central Sonora and the tip of the peninsula were desert gatherers-farmers (Nolasco, 1982). The central desert area of the Baja California peninsula had the least linguistic diversity (and probably the lowest biological diversity; T. Bowen, personal communication).

Loss of biocultural diversity in Mexico (and the Americas in general) began as soon as Europeans arrived. Although estimates of the Native American population size of Mexico upon first arrival of Europeans in 1519 vary greatly (from less than 3 million to over 52 million), most researchers put it at around 20 million (Koch, Brierley, Maslin, & Lewis,



2019). At the time, large areas of land were under cultivation with maize, cacao, and fruit orchards (Whitmore & Turner, 1992). This indigenous population quickly began to collapse from warfare, slavery, and the introduction of pathogens unknown in the Americas (e.g., influenza, smallpox, bubonic plague). The most devastating epidemic in Mexico occurred in 1520, when a single smallpox outbreak killed an estimated 30 to 50 % of the indigenous population (Cook & Borah, 1960; Dobyns, 1993; McCaa, 1995). Following the first comprehensive census in 1568, the population of central Mexico had already declined to 2.7 million (Sanders, Pasons, & Santley, 1979), which corresponds to an approximate decline of 87 % within the first 50 years of European arrival (based on a contact population of 20 million) (Whitmore & Turner, 1992). A rapid population collapse of up to 90 % is plausible for the most populous parts of the country (Koch et al., 2019). One can only speculate how many indigenous languages were lost or put on a path toward extinction during this period. In Northern Mexico many groups of hunters and gatherers who spoke unknown languages perished because of the extermination campaigns launched by Spanish conquerors (Borah & Cook, 1963).

Today, there is strong geographical overlap between the states that harbor the bulk of Mexico's biological and linguistic diversity, with Oaxaca, Puebla, Chiapas, Veracruz, Guerrero and Michoacán standing out. Furthermore, Mexico's biodiversity hotspots closely mirror its language hotspots: areas with the highest number of endangered plant and animal species overlap with those where the endangerment of languages is greatest. Most of Mexico's dry forests, tropical rain forests, and temperate rain forests, which are home to high levels of biological and linguistic diversity, belong to indigenous communities, and almost a third of the country's federally protected areas are within indigenous territories (Sarukhán et al., 2017). It has been estimated that nearly 90 % of Mexico's indigenous population live in forested areas, while the rest live in arid and

semi-arid regions with shrub or grasslands (Toledo et al., 2001a, 2001b).

Oaxaca is a good example of how the geography of languages and biodiversity overlap and have evolved together. Its complex geological history produced an elaborate topography and highly diverse range of ecosystems, from tropical coastal areas to temperate pine-oak forests, thorn scrub and cloud forests, which in turn favored adaptive radiation, speciation, and a high diversity of flora and fauna (Ordoñez, 2004). At least 8 431 species of plants (nearly 40 % of Mexico's known flora) and 4 542 animal species (half of the country's vertebrates and 19 % of its known invertebrates) are present in Oaxaca (García-Mendoza, Ordoñez, & Briones-Salas, 2004; CONABIO, 2008). For more than 10 000 years, its indigenous peoples have dispersed and evolved within the state's diverse ecosystems, and today 157 languages (43 % of Mexico's 364 languages) are spoken in more than 4 000 indigenous communities (de Ávila Blomberg, 2004).

The most insidious threats to Mexico's biodiversity are habitat destruction and fragmentation (mostly by deforestation for agriculture and livestock), overexploitation, invasive species, and climate change. The country has already lost about 70 % of its forest cover, mainly the tropical forests of the Southeast (Sarukhán et al., 2017). Prieto-Amparán et al. (2019) analyzed non-tropical land use in Mexico and found the area of primary forest reduced from 55.8 % in 1990 to 37.7 % in 2017 in their study region (which included temperate primary and secondary forest, human settlements, unvegetated areas, and water bodies). In just one year (2016), 253 000 hectares of Mexico's forests disappeared (Global Forest Watch, 2018). The effects of climate change will exacerbate all those threats (Peterson, Tian, Martínez-Meyer, Soberón, & Sánchez-Cordero, 2005).

The largest areas of forest and wildlands in Mexico are usually communal lands (Stoleson et al., 2005), and at least 70 % of its forested areas are held by ejidos (Segura, 2000; Molnar & White, 2001). However, this is rapidly



changing as ejidos sell their land to private landowners (Stoleson et al., 2005), typically to non-campesinos, and the cultural erosion and loss of traditional knowledge that ensues encourages loss of biodiversity (Laird, 2002).

Languages disappear for a variety of reasons: all speakers may die; speakers may succumb to pressures to speak a different language and forget their mother tongue; speakers may choose to stop speaking their native tongue because of interests in participating in national education, finding jobs that pay salaries and could come with various benefits, or marry into a family with speakers from another language. The threat to Mexico's languages includes a combination of factors such as the reduced number of speakers, their geographical dispersion, predominance of adult speakers, and trends for abandoning transmission of languages to new generations (INALI, 2012). All of this is compounded by a lack of interest or even neglect by authorities, which has led to exclusion of indigenous languages from public and institutional spaces, mass media communications such as radio and television, and their diminishing use among communities and families. Some indigenous people seem to have accepted that their languages will disappear, and they will soon be able to communicate only in Spanish (INALI, 2012), while others desire to reverse the extinction trend. And many more simply do not know what to do to save their (Mexico's) cultural heritage.

Traditional peoples have accumulated vast amounts of ecological knowledge, and that knowledge is embodied in their languages. Thus, as languages go extinct, associated traditional ecological knowledge is also lost (Oviedo et al., 2000). This happens because, in most traditional cultures, knowledge is not recorded in writing but is passed on orally to other groups or new generations. In such cases, the loss of local languages means the loss of the traditional means of knowledge transmission.

The science of biocultural diversity is in its infancy. Although we do not yet fully understand it, there is evidence of ancient and profound connections between biological and

linguistic diversity. When species go extinct entire biological communities and ecosystems can be disrupted or permanently altered. Regardless of their economic, scientific and aesthetic value to humans, plant and animal species have a value of their own because of their unique evolutionary history, genomic diversity, and, ultimately, because of their very existence. When we allow languages to disappear, we squander the culture of humanity, the millennial knowledge of the natural world, and a part of our own past. The IPBES released its definitive new global synthesis of "Assessing nature's contributions to people" (Díaz et al., 2018). It is the first such report since the landmark MEA (2005) was published, and the first ever that is fully intergovernmental. It is also the first assessment to systematically include indigenous and local knowledge at a global scale.

For decades, scientists and conservationists have advocated for the establishment of protected areas to conserve ecosystems and biological diversity (Vidal, López-García, & Rendón-Salinas, 2014). But, as has been advocated for more than two decades, the world's biodiversity will only be effectively preserved by also protecting the diversity of human cultures, and vice versa (Oviedo et al., 2000; Maffi, 2001; Toledo, 2001b). Protecting areas that are implemented under participatory and co-management schemes is the most effective strategy for conserving biodiversity while improving the economic and social conditions of local communities. Conservation area networks comprising priority areas should serve as the baseline for identifying the most suitable strategies depending on the areas' socio-economic contexts and specific characteristics.

We have shown that Mexico's biocultural diversity is at crossroads. Federal, state and municipal governments, businesses, conservation organizations, philanthropists, and multilateral agencies that care about biodiversity need to realize that their resources and efforts will only be effective in the long-term if they simultaneously support protection of indigenous cultures and traditional knowledge. Given



that the regions harboring Mexico's highest biocultural diversity are considered strategic for the country's water, environmental and food security, as well as to ensuring the rights of indigenous peoples, it has recently been proposed that they should be an important component of the 2019-2024 national development plan (Luque & Ortiz Espejel, 2019). We hope this policy initiative results in concrete action to protect Mexico's unique cultural and natural heritage. Biological and cultural diversity are two sides of the same coin. Since endangered languages and endangered species strongly overlap geographically in Mexico, it makes sense to combine efforts to protect both. But time is of the essence. All constituents need to urgently reinforce efforts and significantly augment investments if we are to rescue and preserve the country's unique biocultural diversity and traditional knowledge for the benefit of present and future generations.

Ethical statement: authors declare that they all agree with this publication and made significant contributions; that there is no conflict of interest of any kind; and that we followed all pertinent ethical and legal procedures and requirements. All financial sources are fully and clearly stated in the acknowledgements section. A signed document has been filed in the journal archives.

ACKNOWLEDGMENTS

We are very grateful to linguists Stephen Marlett and Mauricio J. Mixco, to anthropologist Cathy Moser Marlett, and to ethnoecologist Víctor M. Toledo, for their critical reviews of our paper.

RESUMEN

La diversidad biocultural de México en peligro.

Introducción: Lugares con diversidad de especies alta contienen diversidad lingüística alta, mientras que áreas con diversidad de especies baja tienden a contener diversidad lingüística baja. **Objetivo:** Caracterizar las relaciones entre la diversidad biológica y la diversidad cultural, una correlación que ha sido examinada a escala global pero

que en este trabajo es comprobada por primera vez para México. **Métodos:** Recopilamos bases de datos extensas sobre las especies y las lenguas en peligro de extinción, y revisamos la literatura disponible sobre la diversidad biocultural de México, con énfasis en las especies y lenguas en peligro y en peligro crítico de extinción. **Resultados:** Con 364 lenguas vivas, México es el quinto país más diverso lingüísticamente hablando, pero 64 de estas lenguas están en muy alto riesgo de desaparecer y 13 ya desaparecieron. México también es el cuarto país más biodiverso, pero 1 213 especies de su flora y fauna están amenazadas de extinción y al menos 127 desaparecieron recientemente. **Conclusiones:** Los pueblos indígenas son custodios de mucha de la diversidad biocultural del mundo. A medida que el mundo se vuelve menos diverso lingüística y culturalmente, también se vuelve menos diverso biológicamente. La diversidad biológica y lingüística de México muestran una marcada superposición geográfica, y los estados de Oaxaca, Chiapas, Veracruz, Guerrero y Michoacán son los que más especies y lenguajes albergan. De manera similar, los sitios en donde la biodiversidad está en mayor peligro también corresponden con los sitios en donde las lenguas lo están, y las áreas con el mayor número de especies en peligro traslanan con las áreas en donde las lenguas están en mayor peligro.

Palabras clave: idiomas, lingüística, conocimiento tradicional, pueblos indígenas, biodiversidad, México, especies en peligro de extinción, extinción.

REFERENCES

- Berkes, F. (2001). Religious traditions and biodiversity. *Encyclopedia of Biodiversity*, 5, 109-120.
- Beer, N., Gonzalez, S., Huddart, D., Rosales-Lopez, A., & Lamb, A. (2008, May). The palaeodiet of the Pericue Indians of the Cape region of Baja California Sur, Mexico. *Eos Transactions American Geophysical Union*, 89(23), Jt. Assem. Suppl., Abstract U54B-03.
- Birchall, J., Dunn, M., & Greenhill, S.J. (2016). A combined comparative and phylogenetic analysis of the Chapacuran Language family. *International Journal of American Linguistics*, 82(3), 255-284.
- Boas, F. (1917). El dialecto mexicano de Pochutla, Oaxaca. *International Journal of American Linguistics*, 1(1), 9-44. Retrieved from <https://bibliotecatilcoatl.files.wordpress.com/2015/04/el-dialecto-mexicano-de-pochutla-oax-boas-2-37.pdf>
- Boege, E. (2008). *El patrimonio biocultural de los pueblos indígenas de México: hacia la conservación in situ de la biodiversidad y agrodiversidad en los territorios indígenas*. México: Instituto Nacional de Antropología e Historia, Comisión Nacional para el Desarrollo de Pueblos Indígenas.



- Boege, E. (2009). El reto de la conservación de la biodiversidad en los territorios de los pueblos indígenas. In CONABIO, *Capital natural de México* (Vol. II): *Estado de conservación y tendencias de cambio* (pp. 603-649). México: CONABIO.
- Borah, W., & Cook, S.F. (1963). *The aboriginal population of central Mexico on the eve of the Spanish conquest*. Ibero-American Series, No. 45. Berkeley, CA: University of California Press.
- Bouckaert, R., Lemey, P., Dunn, M., Greenhill, S.J., Alekseyenko, A.V., Druimmond, A.J., . . . Atkinson, Q.D. (2012). Mapping the origins and expansion of the Indo-European language family. *Science*, 337, 957-960.
- Bowen, T. (1976). *Seri Prehistory: The Archaeology of the Central Coast of Sonora, Mexico*. Anthropological Papers, No. 27. Tucson, AZ: University of Arizona Press.
- Bowen, T. (2000). *Unknown Island. Seri Indians, Europeans, and San Esteban Island in the Gulf of California*. Albuquerque, NM: University of New Mexico Press.
- Bowen, T. (2006). Recursos culturales de la región de las Grandes Islas en el Golfo de California. *Gaceta Ecológica (Instituto Nacional de Ecología)*, 81, 19-29.
- Bowen, T. (2009). *The record of Native People on Gulf of California Islands*. Arizona State Museum Archaeological Series, No. 201. Tucson, AZ: University of Arizona Press.
- Brusca, R.C., Álvarez-Borrego, S., Hastings, P.A., & Findley, L.T. (2017). Colorado River flow and biological productivity in the Northern Gulf of California, Mexico. *Earth-Science Reviews*, 164, 1-30.
- Brusca, R.C., Moore, W., & Shuster, S.E. (2016). *Invertebrates* (3rd ed.). Sunderland, MA: Sinauer Associates.
- Bryan, J., Shearman, P., Ash, J., & Kirkpatrick, J.B. (2010). Estimating rainforest biomass stocks and carbon loss from deforestation and degradation in Papua New Guinea 1972-2002: Best estimates, uncertainties and research needs. *Journal of Environmental Management*, 91, 995-1001.
- Caccavari Garza, E. (2014). Lenguas Yumanas: crisis de la diversidad lingüística en Baja California. *Revista Digital Universitaria (UNAM)*, 15 (2), 1-12. Retrieved from <http://www.revista.unam.mx/vol.15/num2/art13/art13.pdf>
- Campbell, L. (1997). *American Indian languages: The historical linguistics of Native America*. New York, NY: Oxford University Press.
- Comité International para la Recuperación de la Vaquita (CIRVA). (2019). Report of the Eleventh Meeting of the CIRVA, February 19-21, Southwest Fisheries Science Center. La Jolla, CA: CIRVA.
- Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO). (2008). *Capital natural de México* (Vol. I): *Conocimiento actual de la biodiversidad*. México: CONABIO.
- Comisión Nacional para el Conocimiento y Uso de la Biodiversidad (CONABIO). (2016). *Estrategia nacional sobre la biodiversidad de México y plan de acción 2016-2030*. México: CONABIO.
- Cook, S.F., & Borah, W. (1960). *The Indian population of central Mexico, 1531-1610*. Berkeley, CA: University of California Press.
- Corlett, R.T., & Primack, R.B. (2010). *Tropical rain forests: An ecological and biogeographical comparison*. Malden, MA: Blackwell Publishing.
- Cunningham, A.B. (2001). *Applied ethnobotany: People, wild plant uses and conservation*. London, U.K.: Earthscan.
- Dávila, P., Arizmendi, M.C., Valiente-Banuet, A., Villa- seño, J.L., Casas, A., & Lira, R. (2002). Biological diversity in the Tehuacán-Cuicatlán valley, Mexico. *Biodiversity and Conservation*, 11, 421-442.
- De Ávila Blomberg, A. (2004). La clasificación de la vida en las lenguas de Oaxaca. In A.J. García-Mendoza, M.J. Ordoñez, & M. Briones-Salas (Eds.), *Biodiversidad de Oaxaca* (pp. 481-539). México: Instituto de Biología UNAM, Fondo Oaxaqueño para la Conservación de la Naturaleza, World Wildlife Fund.
- Diario Oficial de la Federación. (2008). Catálogo de las lenguas indígenas nacionales: variantes lingüísticas de México con sus autodenominaciones y referencias geostadísticas. Ciudad de México, México: Instituto Nacional de Lenguas Indígenas (INALI).
- Diario Oficial de la Federación. (2010a). *Programa de revitalización, fortalecimiento y desarrollo de las lenguas indígenas nacionales 2008-2012, PINALI*. Ciudad de México, México: Instituto Nacional de Lenguas Indígenas (INALI).
- Diario Oficial de la Federación. (2010b). *NORMA Oficial Mexicana NOM-059-SEMARNAT-2010, Protección ambiental-Especies nativas de México de flora y fauna silvestres-Categorías de riesgo y especificaciones para su inclusión, exclusión o cambio-Lista de especies en riesgo*. Ciudad de México, México: Secretaría de Medio Ambiente y Recursos Naturales (SEMARNAT).
- Díaz, S., Pascual, U., Stenseke, M., Martín-López, B., Watson, R.T., Molnár, Z., . . . Shirayama, Y. (2018). Assessing nature's contributions to people. *Science*, 359(6373), 270-272.



- Dobyns, H.F. (1993). Disease transfer at contact. *Annual Reviews of Anthropology*, 22, 273-291.
- Dunn, M. (2014). Language phylogenies. In C. Bowern, & B. Evans (Eds.), *Routledge Handbook of Historical Linguistics* (pp. 190-211). New York, NY: Routledge.
- Felger, R.S., & Moser, M.B. (1987). Sea turtles in Seri Indian culture. *Environment Southwest, Autumn*, 18-21.
- Felger, R.S., Nichols, W.J., & Seminoff, J.A. (2005). Sea turtles in northwestern Mexico: Conservation, ethno-biology, and desperation. In J.E. Cartron, G. Ceballos, & R.S. Felger (Eds.), *Biodiversity, Ecosystems, and Conservation in Northern Mexico* (pp. 405-424). New York, NY: Oxford University Press.
- Fincher, C.L., & Thornhill, R. (2008). A parasite-driven wedge: Infectious diseases may explain language and other biodiversity. *Oikos*, 117, 1289-1297.
- Fujita, H. (2006). The Cape region. In D. Laylander & J.D. Moore (Eds.), *The Prehistory of Baja California: Advances in the Archaeology of the Forgotten Peninsula* (pp. 82-98). Gainesville, FL: University Press of Florida.
- García-Mendoza, A.J., Ordoñez, M.J., & Briones-Salas, M. (Eds.). (2004). *Biodiversidad de Oaxaca*. Ciudad de México, México: Universidad Nacional Autónoma de México.
- Gavin, M.C., Botero, C.A., Bowern, C., Colwell, R.K., Dunn, M., Dunn, R.R., ... Yanega, G. (2013). Toward a mechanistic understanding of linguistic diversity. *BioScience*, 63, 524-535.
- Gavin, M.C., McCarter, J., Mead, A., Berkes, F., Stepp, J.R., Peterson, D., & Tang, R. (2015). Defining biocultural approaches to conservation. *Trends in Ecology & Evolution*, 30(3), 140-145.
- Global Forest Watch [Website]. (2018). Retrieved from <https://www.globalforestwatch.org/>
- Gonzalez-Pérez, G., Briones-Salas, M., & Alfaro, A.M. (2004). Integración del conocimiento faunístico del estado. In A.J. García-Mendoza, M.J. Ordoñez, & M. Briones-Salas (Eds.), *Biodiversidad de Oaxaca* (pp. 449-466). México: Instituto de Biología UNAM, Fondo Oaxaqueño para la Conservación de la Naturaleza, World Wildlife Fund.
- Gorenflo, L.J., Romaine, S., Mittermeier, R.A., & Walker-Painemilla, K. (2012). Co-occurrence of linguistic and biological diversity in biodiversity hotspots and high biodiversity wilderness areas. *Proceedings of the National Academy of Sciences*, 109(21), 8032-8037.
- Gray, R.D., Quentin, D., & Greenhill, S.J. (2018). Language evolution and human history: What a difference a date makes. *Philosophical Transactions of The Royal Society B: Biological Sciences*, 366(1567), 1090-1100.
- Hamilton, M.J., Milne, B.T., Walker, R.S., Burger, O., & Brown, J.H. (2007). The complex structure of hunter-gatherer social networks. *Proceedings of the Royal Society B: Biological Sciences*, 274(1622), 2195-2203.
- Hammarström, H., Forkel, R., & Haspelmath, M. (Eds.). (2018). Glottolog 3.3 [Software]. Available from <http://glottolog.org>
- Harmon, D. (1995). The status of the world's languages as reported in 'Ethnologue.' *Southwest Journal of Linguistics*, 14, 1-28.
- Harmon, D. (1996). Losing species, losing languages: Connections between linguistic and biological diversity. *Southwest Journal of Linguistics*, 15, 89-108.
- Harmon, D., & Loh, J. (2010). The index of linguistic diversity: A new quantitative measure of trends in the status of the world's languages. *Language, Documentation and Conservation*, 4, 97-151.
- Harrison, K.D. (2007). *When languages die: The extinction of the world's languages and the erosion of human knowledge*. London, U.K.: Oxford University Press.
- Holt, D. (2001). Valedictory: Lino de la Rosa. *Ogmios Newsletter of Foundation for Endangered Languages* 2.4, 16, 30. Retrieved from <https://www.ogmios.org/ogmios>
- Instituto Nacional de Lenguas Indígenas (INALI). (2010). *Proyecto de documentación inicial de la lengua ku'ahl* (Informe Ejecutivo, Contrato INALI.C.B.17/020/2010). México: INALI. Retrieved from http://site.inali.gob.mx/pdf/estudios_opiniones/Documentacion_kuahl.pdf
- Instituto Nacional de Lenguas Indígenas (INALI). (2012). *Lenguas indígenas nacionales en riesgo de desaparición: Variantes lingüísticas por grado de riesgo*. A. Embriz Osorio & O. Zamora Alarcón (Coords.). México: INALI.
- Instituto Nacional de Estadística y Geografía (INEGI). (2010). *Censo de población y vivienda 2010*. México: INEGI. Retrieved from <https://www.beta.inegi.org.mx/proyectos/ccpv/2010/>
- Instituto Nacional de Estadística y Geografía (INEGI). (2015). *Resultados de la encuesta intercensal 2015. Tabulados básicos*. México: INEGI. Retrieved from <https://www.inegi.org.mx/est/contenidos/Proyectos/encuestas/hogares/especiales/ ei2015/>
- Inter-American Commission on Human Rights. (2009). *Indigenous and tribal peoples' rights over their ancestral lands and natural resources* (OEA/Ser.L/V/II, Doc. 56/09, 30 December 2009). Washington, DC: OAS. Retrieved from <http://cidh.org/countryrep/Indigenous-Lands09/TOC.htm>



- International Union for the Conservation of Nature (IUCN). (2018). The IUCN Red List of Threatened Species. Version 2018-2 [Website]. Retrieved October 09, 2018 from <https://www.iucnredlist.org/>
- Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services (IPBES). (2018). *The IPBES regional assessment report on biodiversity and ecosystem services for the Americas*. J. Rice, C.S. Seixas, M.E. Zaccagnini, M. Bedoya-Gaitán, & N. Valderrama (Eds.). Bonn, Germany: Secretariat of the IPBES.
- Jones, K.R., Venter, O.P., Fuller, R.A., Allan, J.R., Maxwell, S.L., Negret, P.J., & Watson, J.E.M. (2018). One-third of global protected land is under intense human pressure. *Science*, 360(6390), 788-791.
- Kerr, J.T., & Packer, L. (1997). Habitat heterogeneity as a determinant of mammal species richness. *Nature*, 385(6613), 252-254.
- Knab, T. (1980). When is a language really dead: The case of Pochutec. *International Journal of American Linguistics*, 46(3), 230-233. <https://doi.org/10.1086/465658>
- Koch, A., Brierley, C., Maslin, M.M., & Lewis, S.L. (2019). Earth system impacts of the European arrival and great dying in the Americas after 1492. *Quaternary Science Reviews*, 207, 13-36.
- Laird, S.A. (Ed.). (2002). *Biodiversity and traditional knowledge: Equitable partnerships in practice*. London, U.K.: Earthscan Publications.
- Lewis, M., Simons, P.G.F., & Fennig, C.D. (2016). *Ethnologue: Languages of the world* (19th ed.). Dallas, TX: SIL International.
- Lionnet, A. (1978). *El idioma Tubar y los Tubares*. Ciudad de México, México: Universidad Iberoamericana.
- Lionnet, A. (1986). *El Eudeve: Un idioma extinto de Sonora*. Ciudad de México, México: Universidad Nacional Autónoma de México.
- Loh, J., & Harmon, D. (2005). A global index of biocultural diversity. *Ecological Indicators*, 5, 231-241.
- Loh, J., & Harmon, D. (2014). *Biocultural diversity: Threatened species, endangered languages*. Zeist, The Netherlands: WWF Netherlands.
- Luque, D., & Ortiz Espejel, B. (2019). *Hacia una política de bienestar comunitario de pueblos indígenas y comunidades equiparables en regiones de alta densidad biocultural de México* (Propuesta elaborada por la Red Temática CONACYT Patrimonio Biocultural de México). México: CONACYT.
- Maffi, L. (1998). Language: A resource for nature. *Nature and Resources: The UNESCO Journal on the Environment and Natural Resources Research*, 34, 12-21.
- Maffi, L. (Ed.). (2001). *On biocultural diversity: Linking language, knowledge and the environment*. Washington, DC: Smithsonian Institution Press.
- Maffi, L. (2005). Linguistic, cultural, and biological diversity. *Annual Reviews of Anthropology*, 29, 599-617.
- Marlett, C.M. (2014). *Shells on a desert shore: Mollusks in the Seri world*. Tucson, AZ: University of Arizona Press.
- McCaa, R. (1995). Spanish and Nahuatl views on smallpox and demographic catastrophe in Mexico. *Journal of Interdisciplinary History*, 25, 397-431.
- Millennium Ecosystem Assessment (MEA). (2005). *Ecosystems and human well-being: Synthesis*. Washington, DC: Island Press.
- Mittermeier, R.A., & Mittermeier, C.G. (1992). La importancia de la diversidad biológica de México. In J. Sarukhán & R. Dirzo (Eds.), *México ante los retos de la biodiversidad* (pp. 6373). Ciudad de México, México: CONABIO.
- Mittermeier, R.A., Myers, N., Gil, P.R., & Mittermeier, C.G. 1999. *Hotspots: Earth's biologically richest and most endangered terrestrial ecoregions*. Monterrey, México: Cemex, Conservation International, Agrupación Sierra Madre.
- Mixco, M.J. (1978). *Cochimí and Proto-Yuman: Lexical and syntactic evidence for a new language family in Lower California*. University of Utah Anthropological Papers 101. Salt Lake City, UT: University of Utah Press.
- Molina Cruz, M. (2010). *La recuperación de la lengua Xuaní-Ixcateca de Oaxaca, a través del video* (Tesis de Licenciatura). Universidad Pedagógica Nacional Unidad Ajusco, México.
- Molnar, A., & White, T.A. (2001). Forestry and land management. In M.M. Giugale, O. Lafourcade, & V.H. Nguyen (Eds.), *Mexico: A comprehensive development agenda for the new era* (pp. 669-690). Washington, DC: World Bank.
- Moore, J.L., Manne, L., Brooks, T., Burgess, N.D., Davies, R., Rahbek, C., . . . Balmford, A. (2002). The distribution of cultural and biological diversity in Africa. *Proceedings of the Royal Society B: Biological Sciences*, 269(1501), 1645-1653.
- Mora, C., Tittensor, D.P., Adl, S., Simpson, A.G.B., & Worm, B. (2011). How many species are there on Earth and in the ocean? *PLoS Biology*, 9(8), e1001127. <https://doi.org/10.1371/journal.pbio.1001127>
- Mühlhäuser, P. (1995). *The Politics of Multiculturalism in the Asia/Pacific*. Darwin, Australia: Northern Territory University Press.



- Raygorodetsky, G. (2018). Indigenous peoples defend Earth's biodiversity—but they're in danger. Retrieved November 17, 2018, from <https://www.nationalgeographic.com/environment/2018/11/can-indigenous-land-stewardship-protect-biodiversity/>
- Nettle, D. (1996). Language diversity in West Africa: An ecological approach. *Journal of Anthropologic Archaeology*, 15, 403-438.
- Nettle, D. (1999a). Is the rate of linguistic change constant? *Lingua*, 108, 119-136.
- Nettle, D. (1999b). *Linguistic diversity*. New York, NY: Oxford University Press.
- Nettle, D., & Romaine, S. (2000). *Vanishing Voices*. New York, NY: Oxford University Press.
- Nolasco Armas, M. (1982). De nómadas cazadores y recolectores a aldeas agrícolas indiferenciadas (el caso del noroeste de México). *Anales de Antropología*, 19, 59-72.
- Ordoñez, M.J. (2004). El territorio. In A.J. García-Mendoza, M.J. Ordoñez, & M. Briones-Salas (Eds.), *Biodiversidad de Oaxaca* (pp. 469-479). México: Instituto de Biología UNAM, Fondo Oaxaqueño para la Conservación de la Naturaleza, World Wildlife Fund.
- Oviedo, G., Maffi, L., & Larsen, P.B. (2000). *Indigenous and traditional peoples of the world ecoregion conservation. An integrated approach to conserving the world's biological and cultural diversity*. Gland, Switzerland: World Wide Fund for Nature International.
- Papua New Guinea's Fifth National Report to the Convention on Biological Diversity*. (2014). Port Moresby, Papua New Guinea: Government of Papua New Guinea.
- Perales, H.R., & Aguirre, J.R. (2008). Biodiversidad humanaizada. In CONABIO, *Capital Natural de México* (Vol. I): *Conocimiento actual de la biodiversidad* (pp. 565-603). México: CONABIO.
- Peterson A.T., Tian, H., Martínez-Meyer, E., Soberón, J., & Sánchez-Cordero, V. (2005). Modeling ecosystems shifts and individual species distribution shifts. In T. Lovejoy & L. Hannah (Eds.), *Climate change and biodiversity* (pp. 45-78). Washington, DC: Smithsonian Institution Press.
- Pōmaika'i McGregor, D. (1999). Research in action: Ethnohistory of Puna. *Social Process in Hawaii*. 39, 181-207.
- Posey, D.A. (1999). Cultural and spiritual values of biodiversity. A complementary contribution to the global biodiversity assessment. In D.A. Posey (Ed.), *Cultural and Spiritual Values of Biodiversity* (pp. 1-19). London, U.K: UNEP and Intermediate Technology Publications.
- Prieto-Amparán, J.A., Villarreal-Guerrero, F., Martínez-Salvador, M., Manjarrez-Domínguez, C., Vázquez-Quintero, G., & Pinedo-Álvarez, A. (2019). Spatial near future modeling of land use and land cover changes in the temperate forests of Mexico. *PeerJ Preprints*, 7, e27542v1 <https://doi.org/10.7287/peerj.preprints.27542v1>.
- Primack, R.C., & Vidal, O. (2019). *Introducción a la biología de la conservación*. Ciudad de México, México: Fondo de Cultura Económica, Ediciones Científicas Universitarias.
- Renfrew, C. (1994). World linguistic diversity. *Scientific American*, 270, 116-123.
- Rojas-Bracho, L., Brusca, R.C., Álvarez-Borrego, S., Brownell Jr., R.L., Camacho-Ibar, V., Ceballos, G., . . . Taylor, B.L. (2018). Unsubstantiated scientific claims can lead to tragic conservation outcomes. *BioScience*, 69(1), 12-14.
- Sanders, W.T., Pasons, J.R., & Santley, R.S. (1979). *The basin of Mexico: The ecological processes in the evolution of a civilization*. New York, NY: Academic Press.
- Sarukhán, J., Koleff, P., Carabias, J., Soberón, J., Dirzo, R., Llorente-Bousquets, J., . . . García Méndez, G. (2017). *Capital natural de México. Síntesis: evaluación del conocimiento y tendencias de cambio, perspectivas de sustentabilidad, capacidades humanas e institucionales*. México: CONABIO.
- Segura, G. (2000, May). Mexico's forest sector and policies: A general perspective. *Constituting the Commons: Crafting Sustainable Commons in the New Millennium*. Paper presented at the International Association for the Study of Common Property (IASCP) Biennial Conference, Bloomington, IN.
- Shengji, P. (1999). The holy hills of the Dai. In D.A. Posey (Ed.), *Cultural and Spiritual Values of Biodiversity* (pp. 381-382). London, U.K.: UNEP and Intermediate Technology Publications.
- Simons, G.F., & Fennig, C.D. (Eds.). (2018). *Ethnologue: languages of the world* (21st ed.). Dallas, TX: SIL International. Retrieved from <https://www.ethnologue.com/ethnoblog/gary-simons/welcome-21st-edition>
- Stepp, J.R., Cervone, S., Castaneda, H., Lasseter, A., Stocks, G., & Gichon, Y. (2004). Development of a GIS for global biocultural diversity. *Policy Matters*, 13, 267-270.
- Stoleson, S.H., Felger, R.S., Ceballos, G., Raish, C., Wilson, M.F., & Búrquez, A. (2005). Recent history of natural resource and population growth in northern Mexico. In J.E. Carrion, G. Ceballos, & R.S. Felger (Eds.), *Biodiversity, Ecosystems, and Conservation in*



- Northern Mexico (pp. 52-86). New York, NY: Oxford University Press.
- Sutherland, W.J. (2003). Parallel extinction risk and global distribution of languages and species. *Nature*, 423, 276-279.
- Thomas, L., Jaramillo-Legorreta, A., Cárdenas-Hinojosa, G., Nieto-García, E., Rojas-Bracho, L., Ver Hoef, J.M., . . . Tregenza, N. (2017). Last call: Passive acoustic monitoring shows continued rapid decline of critically endangered vaquita. *Journal of the Acoustic Society of America*, 142(5), EL512-EL517.
- Toledo, V.M. (1994). Biodiversity and cultural diversity in Mexico. *Different Drummer*, 1(3), 16-19.
- Toledo, V.M. (2001a). Biocultural diversity and local power in Mexico. In L. Maffi (Ed.), *On biocultural diversity: Linking language, knowledge and the environment* (pp. 472-488). Washington, DC: Smithsonian Institution Press.
- Toledo, V.M. (2001b). Indigenous peoples and biodiversity. In S.A. Levin (Ed.), *Encyclopedia of Biodiversity* (pp. 451-463). San Diego, CA: Academic Press.
- Toledo, V.M., Alarcón-Chaires, P., Moguel, P., Olivo, M., Cabrera, A., Leyequien, E., & Rodríguez-Aldabe, A. (2001). El atlas etnoecológico de México y Centroamérica: fundamentos, métodos y resultados. *Etnoecológica*, 6(8), 7-41.
- Toledo, V.M., & Barrera-Bassols, N. (2008). *La memoria biocultural: La importancia ecológica de las sabidurías tradicionales*. Barcelona, Spain: Icaria Editorial, S.A.
- Toledo, V.M., Boege, E., & Carrera-Bassols, N. (2010). The Biocultural heritage of Mexico: An overview. *Landscape*, 2(6), 7-13.
- United Nations Educational, Scientific and Cultural Organization (UNESCO). (2010). *Atlas of the world's languages in danger*. Paris, France: UNESCO.
- Vidal, O. (1995). Population biology and incidental mortality of the vaquita, *Phocoena sinus*. *Reports of the International Whaling Commission, Special Issue*, 16, 247-272.
- Vidal, O., López-García, J., & Rendón-Salinas, E. (2014). Trends in deforestation and forest degradation after a decade of monitoring in the monarch butterfly biosphere reserve in Mexico. *Conservation Biology*, 28, 177-186.
- Villaseñor, J.L. (2016). Checklist of the native vascular plants of Mexico. *Revista Mexicana de Biodiversidad*, 84, 559-902.
- Whitmore, T.M., & Turner, B.L. (1992). Landscapes of cultivation in Mesoamerica on the eve of the conquest. *Annals of the Association of American Geographers*, 82, 402-425.
- Wiersum, K.F. (2004). Use and conservation of biodiversity in East African forested landscapes. In P.A. Zuidema (Ed.), *Tropical forests in multi-functional landscapes. Proceedings of Seminar Series Issues in International Nature Conservation* (pp. 33-39). Utrecht, The Netherlands: Utrecht University Prince Bernard Centre for International Nature Conservation.



SUPPLEMENTAL MATERIAL

Common and scientific names of endemic and non-endemic critically endangered and endangered species from Mexico (Diario Oficial de la Federación 2010)

FUNGI

Tricholosporum subporphyrophyllum, non-endemic
Tricholosporum tropicalis, non-endemic
Conocybe siligineoides, non-endemic
Hypholoma naematoliformis, non-endemic
Psilocybe heimii, non-endemic
Psilocybe pleurocystidiosa, non-endemic
Psilocybe singeri, non-endemic
Psilocybe uxpanapensis, non-endemic
Psilocybe verae-crucis, non-endemic
Psilocybe weldenii, non-endemic

PLANTS

flecha de agua, *Sagittaria intermedia*, non-endemic
Tauschia allioides, non-endemic
camedor despeinado, *Chamaedorea glaucifolia*, endemic
camedor metálico, *Chamaedorea metallica*, endemic
camedor guayita, *Chamaedorea tenella*, endemic
camedor guonay, *Chamaedorea tuerckheimii*, non-endemic
falso camedor, *Synechanthus fibrosus*, non-endemic
maguey de la luna, *Agave lurida*, endemic
maguey de Nizanda, *Agave nizandensis*, endemic
Agave victoriae-reginae, endemic
Hymenocallis concinna, endemic
Hymenocallis durangoensis, endemic
Petronympha decora, endemic
soyate de Purpus, *Beaucarnea purpusii*, endemic
encyclia de Kienast, *Encyclia kienastii*, endemic
galeottia grande, *Galeottia grandiflora*, non-endemic
laelia de Muertos, lirios, *Laelia anceps dawsonii*, endemic
lycaste pelosa, *Lycaste lassiglossa*, non-endemic
lycaste monjita, *Lycaste skinneri*, non-endemic
Mexipedium xerophyticum, endemic
mormodes sanguíneo, *Mormodes sanguineoclastra*, endemic
mormodes de Soto, *Mormodes sotoana*, non-endemic
mormodes fimbriado, *Mormodes uncia*, endemic
tanal de bigotes, *Phragmipedium extaminodium*, non-endemic
odontoglossum de mayo, *Rhynchostele majalis*, non-endemic
Rhynchostele uroskinneri, non-endemic
odontoglossum grande, *Rossioglossum grande*, non-endemic
odontoglossum de Williams, *Rossioglossum williamsianum*, non-endemic
trichopilia amarilla, *Trichopilia galeottiana*, non-endemic
girasol, *Hazardia orcuttii*
Perymenium wilburorum, non-endemic
Villasenorbia orcuttii, non-endemic
cactus junco floricuelo, *Aporocactus flagelliformis*, endemic
biznaga peyotillo, *Ariocarpus fissuratus bravoanus*, endemic
Ariocarpus scaphirostris, endemic
biznaga algodoncillo de estrella, cacto estrella, *Astrophytum asterias*, endemic

biznaga partida delgada, *Coryphantha gracilis*, endemic
biznaga partida amacollada, *Coryphantha werdermannii*, endemic
Digitostigma caput-medusae, endemic
biznaga tonel dorada, *Echinocactus grusonii*, endemic
órgano pequeño de Jaraguay, *Echinocereus ferreirianus lindsayi*, endemic
órgano pequeño de Querétaro, *Echinocereus schmollii*, endemic
Echinomastus erectocentrus acunensis, non-endemic
Lophophora diffusa viridiscens, endemic
biznaga de la Reja, *Mammillaria carmenae*, endemic
tezontle, *Mammillaria crinita leucanth*, endemic
biznaga pol tsakam, *Mammillaria gaumeri*, endemic
biznaga de San Ángel, *Mammillaria haageana san-angelensis*, endemic
biznaga bola de hilo, *Mammillaria herrerae*, endemic
biznaga de Lau, *Mammillaria laui dasyacantha*, endemic
biznaga de Lau, *Mammillaria laui laui*, endemic
biznaga de La Cañada, *Mammillaria mathilda*, endemic
Mammillaria sanchez-mejoradae, endemic
biznaga de Coneto, *Mammillaria theresae*, endemic
Melocactus curvispinus curvispinus, non-endemic
Pterocereus gaumeri, endemic
biznaga cono invertido de Gielsdorf, *Turbinicarpus gielsdorffianus*, endemic
Turbinicarpus jauernigii, endemic
turbinita de Querétaro, *Turbinicarpus pseudomacrochele*, endemic
uñita, *Turbinicarpus schmiedickeanus andersonii*, endemic
uñita, *Turbinicarpus schmiedickeanus gracilis*, endemic
Turbinicarpus schmiedickeanus rioverdensis, endemic
Zinowiewia concinna, non-endemic
Cyathea costaricensis, non-endemic
Nephelea mexicana, non-endemic
palmita (Chiapas), *Ceratozamia alvarezii*, endemic
mazacopa (Zoque), *Ceratozamia chimalapensis*, endemic
palmita (Veracruz), *Ceratozamia decumbens*, endemic
Ceratozamia kuesteriana, endemic
Ceratozamia latifolia, endemic
Ceratozamia matudae, endemic
palmita, *Ceratozamia miqueliana*, endemic
amendaui (Zoque), *Ceratozamia mirandae*, endemic
carrete (Oaxaca), *Ceratozamia mixeorum*, endemic
tepemaizte (Náhuatl, Veracruz), *Ceratozamia morettii tepetmaizte*, endemic
Ceratozamia norstogii, endemic
palma, palma espinosa, amendu, espadaña (Chiapas), *Ceratozamia vovidesii amenduai*, endemic
Ceratozamia whitelockiana, endemic
Ceratozamia zaragozae, endemic
Ceratozamia zoquorum, endemic
chamal (Nuevo León, Tamaulipas), *Dioon angustifolium*, endemic
palma (Oaxaca), *Dioon argenteum*, endemic
palma (Oaxaca), *Dioon califanoi*, endemic
palma real (Oaxaca), *Dioon caputoi*, endemic
chamal (Nuevo León), palma de Teresita (Tamaulipas), palma de dolores (San Luis Potosí), palma navaja (Querétaro), quiotamal, tiotamal (Veracruz), *Dioon edule*, endemic
Marisol, plumilla (Oaxaca), *Dioon holmgrenii*, endemic
Espadaña, nimalari (Chiapas), *Dioon merolae*, endemic
palma real (Oaxaca), *Dioon purpusii*, endemic
tush-kju (Mazateco), *Dioon rzedowskii*, endemic
palma de la Virgen (Sonora, Sinaloa), peyote (Sonora), *Dioon sonorensis*, endemic
palma de chicalite, palma de Dolores, chicalitos, coyolillo, coyolito de cerro (Oaxaca), *Dioon spinulosum*,

endemic
palma de la Virgen (Sinaloa, Durango), palma, palmita (Guerrero, Michoacán, Nayarit), *Dioon tomasellii*, endémica
Zamia cremnophila, endemic
chamalillo (Querétaro, San Luis Potosí), *Zamia fischeri*, endemic
palma bola (Veracruz), *Zamia furfuracea*, endemic
palmita (Veracruz), *Zamia inermis*, endemic
Zamia katzeriana, endemic
Zamia lacandona, endemic
Zamia purpurea, endemic
Zamia sonorensis, endemic
Zamia spartea, endemic
amigo del maíz (Veracruz), *Zamia vazquezii*, endemic
Dicksonia regalis, non-endemic
Dicksonia schiedei, non-endemic
Sideroxylon cartilagineum, non-endemic
Diospyros riojae, non-endemic
zapote prieto, *Diospyros xolocotzii*, endemic
Fouquieria ochoterenae, endemic
Fouquieria purpusii, endemic
Dalbergia congestiflora, non-endemic
Dalbergia granadillo, non-endemic
Ormosia isthmensis, non-endemic
Ormosia macrocalyx, non-endemic
Platymiscium lasiocarpum, non-endemic
Vatarea lundellii, non-endemic
guichín, acailite, pepinque, *Fagus grandifolia* var. Mexicana, endemic
menta espinosa de San Diego, *Acanthomintha ilicifolia*, non-endemic
Litsea glaucescens, non-endemic
Eichhornia azurea, non-endemic
Eichhornia heterosperma, non-endemic
Eichhornia paniculata, non-endemic
Eurystemon mexicanum, non-endemic
Heteranthera oblongifolia, non-endemic
Heteranthera seubertiae, non-endemic
Magnolia dealbata, non-endemic
Euphorbia confertii, endemic
lomboi de playa, *Jatropha giffordiana*, endemic
fremontia, *Fremontodendron mexicanum*, non-endemic
Mortoniodendron guatemalense, non-endemic
Tilia americana mexicana, non-endemic
Potamogeton amplifolius, non-endemic
Potamogeton praelongus, non-endemic
Nymphaea novogranatensis, non-endemic
ciprés de Tecate, ciprés negro, *Cupressus forbesii*, endemic
ciprés brillante, ciprés de Guadalupe, *Cupressus guadalupensis*, endemic
oyamel de Jalisco, *Abies guatemalensis*, non-endemic
oyamel de Juárez, *Abies hickelii*, endemic
pinabete espinoso, *Picea chihuahuana*, non-endemic
Picea engelmannii mexicana, non-endemic
pinabete de Nuevo León, *Picea martinezii*, non-endemic
Pinus attenuata, non-endemic
pino de Honduras, *Pinus caribaea hondurensis*, non-endemic
pino de brea, *Pinus coulteri*, non-endemic
piñón de octubre, *Pinus culminicola*, endemic
pino de Jalisco, *Pinus jaliscana*, endemic

pino azul, *Pinus maximartinezii*, endemic
pino peninsular, *Pinus muricata*, non-endemic
piñón de Nelson, *Pinus nelsonii*, endemic
pino piñonero llorón, *Pinus pinceana*, endemic
pino de Coalcomán, *Pinus rzedowskii*, endemic
Guadua spinosa, non-endemic
Olmeca recta, endemic
Olmeca reflexa, endemic
Orcuttia californica, non-endemic
Triniochloa laxa, non-endemic
Triniochloa micrantha, non-endemic
Zea perennis, endemic
Oserya longifolia, endemic
Nephrolepis cordifolia, non-endemic
Hydrangea nebulicola, endemic
Arce, maple, *Acer saccharum skutchii*, non-endemic
Echeveria elegans, endemic
Echeveria laui, endemic
Echeveria purpusorum, endemic
Echeveria setosa ciliata, endemic
Echeveria setosa deminuta, endemic
Echeveria setosa minor, endemic
Echeveria setosa oteroii, endemic
Echeveria setosa setosa, endemic
Graptopetalum macdougallii, endemic
Sedum frutescens, endemic
Sedum suaveolens, endemic
Louteridium donnell-smithii, non-endemic
Hesperalaea palmeri, non-endemic
Selaginella porphyrospora, non-endemic
Sparganium americanum, non-endemic
Sparganium eurycarpum, non-endemic
huevos de víbora, *Amoreuxia wrightii*, non-endemic
Frankenia johnstonii, non-endemic

INVERTEBRATES

tábano de las dunas, *Brennania belkini*
camaroncillo, *Potamalpheops stygicola*, endemic
chacales, *Typhlatya campecheae*
acocil, cangrejo de agua dulce, *Procamarbarus regiomontanus*, endemic
langostino, *Macrobrachium acerontium*
langostino, *Neopalaemon nahuatlus*
langostino, *Troglobimexicanus perezfarfanteae*
cangrejo de la barrancas, *Pseudothelphusa dugesi*, endemic
cangrejo, *Typhlopseudothelphusa mocinoi*
cacerolita de mar, *Limulus polyphemus*
remipedo, *Speleonectes tulumensis*, endemic
Cyrtonaias tampicoensis tecomatenensis
Megalonaia nicklineana
Coahuilix hubbsi, endemic
Cochliopina milleri, endemic
Durangonella coahuilae, endemic
Mexipyrgus churinceanus, endemic
Mexithauma quadripaludium, endemic
Nymphophilus minckleyi, endemic

Paludiscala caramba, endemic

FISH

charal La Caldera, *Menidia bartoni*, endemic
charal tarasco, *Menidia charari*, endemic
charal del Santiago, *Menidia riojai*, endemic
matalote de Sonora, *Catostomus insignis*, non-endemic
matalote jorobado, *Xyrauchen texanus*, endemic
pupo del Lerma, *Algansea barbata*, endemic
carpita La Concha, sardinita Nazas, *Cyprinella alvarezdelvillari*, endemic
sardinita or carpa bocagrande, *Cyprinella bocagrande*, endemic
sardinita or carpa del Cochos, *Cyprinella panarcys*, endemic
sardinita or carpa de Cuatrocíenegas, *Cyprinella xanthicara*, endemic
carpa diabla, *Dionda diaboli*, endemic
carpa del Bravo, *Dionda episcopa*, endemic
carpa quijarona, *Dionda mandibularis*, endemic
carpa manchada, *Dionda melanops*, endemic
carpa manchada, *Dionda melanops*, endemic
carpita del Gila, *Gila intermedia*, non-endemic
carpa de Saltillo, *Gila modesta*, endemic
carpita yaqui, *Gila purpurea*, non-endemic
carpa del Salado, *Notropis saladonis*, endemic
carpa de Parras, *Stypondon signifer*, endemic
almirante, *Millerichthys robustus*, endemic
cachorro de Medialuna, *Cualac tessellatus*, endemic
cachorro Cecilia, *Cyprinodon ceciliae*, endemic
cachorro escondido, *Cyprinodon esconditus*, endemic
cachorro Carbonera, *Cyprinodon fontinalis*, endemic
cachorro de Julimes, burrito de Julimes, *Cyprinodon julimes*, endemic
cachorro cangrejero, *Cyprinodon labiosus*, endemic
cachorro Charco Palmas, *Cyprinodon longidorsalis*, endemic
cachorro escamudo, *Cyprinodon macrolepis*, endemic
cachorro del desierto, *Cyprinodon macularius*, endemic
cachorro gigante, *Cyprinodon maya*, endemic
cachorro del Mezquital, *Cyprinodon meeki*, endemic
cachorro cabezón, *Cyprinodon pachycephalus*, endemic
cachorro boxeador, *Cyprinodon simus*, endemic
cachorro besucón, *Cyprinodon suavium*, endemic
cachorro dorsal larga, *Cyprinodon verecundus*
cachorro Verónica, *Cyprinodon veronicae*
sardinilla de Península, *Fundulus lima*, endemic
sardinita Cuatrocíenegas, *Lucania interioris*, endemic
mexcalpique de Tuxpan, *Allodontichthys hubbsi*, endemic
mexcalpique de escama, *Allodontichthys polylepis*, endemic
mexcalpique de Tamazula, *Allodontichthys tamazulae*, endemic
tiro Catarina, *Allotoca catarinae*, endemic
chorumo, *Allotoca diazi*, endemic
tiro, *Allotoca dugesii*, endemic
tiro rayado, *Allotoca goslinei*, endemic
tiro rayado, *Allotoca regalis*, endemic
mexcalpique cola azul, *Ameca splendens*, endemic
mexcalpique del Toboso, *Characodon audax*, endemic
mexcalpique arcoíris, *Characodon lateralis*, endemic
mexcalpique, *Girardinichthys viviparus*, endemic
mexcalpique michoacano, cherehuita, *Hubbsina turneri*, endemic

tiro rayado, *Skiffia bilineata*, endemic
mexcalpique viejo, *Xenoophorus captivus*, endemic
mexcalpique negro, *Xenotoca melanosoma*, endemic
picote tequila, *Zoogoneticus tequila*, endemic
guayacón San Gregorio, *Gambusia alvarezi*, endemic
guayacón amarillo or del oeste, *Gambusia speciosa*, endemic
topote del Tamesí, *Poecilia latipunctata*, endemic
topote de Teapa, *Poecilia sulphuraria*, endemic
guayacón bonito, *Priapella bonita*, endemic
espada de Monterrey, *Xiphophorus couchianus*, endemic
espada Cuatrociénegas, *Xiphophorus gordoni*, endemic
espada de Múzquiz, *Xiphophorus meyeri*, endemic
espada de Catemaco, *Xiphophorus milleri*, endemic
escamudo de San Cristóbal, *Profundulus hildebrandi*, endemic
espinocho, *Gasterosteus aculeatus microcephalus*
totoaba, *Totoaba macdonaldi*, endemic
dama blanca ciega, *Typhliasina pearsei*, endemic
mojarra caracolera, *Herichthys bartoni*, endemic
mojarra Cuatrociénegas, *Herichthys minckleyi*, endemic
mojarra ojo frío, *Herichthys steindachneri*, non-endemic
cucharita de baja, *Gobiesox juniperoserrai*, non-endemic
perca del Conchos, *Etheostoma australe*, endemic
perca Cuatro Ciénegas, *Etheostoma lugoi*, endemic
lamprea de Jacona, lamprea de Cuitzeo, *Tetrapleurodon geminis*, endemic
lamprea de Chapala, *Tetrapleurodon spadicea*, endemic
bagre ciego duende, *Prietella lundbergi*, non-endemic
bagre ciego de Múzquiz, *Prietella phreatophila*, endemic
bagre de Chiapas, *Lacantunia enigmática*, endemic
anguila ciega yucateca, *Ophisternon infernale*, endemic

REPTILES

lagarto alicante de Bogert, *Abronia bogerti*, endemic
lagarto alicante de Chiszar, *Abronia chiszari*, endemic
lagarto alicante verde, *Abronia ochoterenai*, endemic
lagarto alicante de Cerro Baúl, *Abronia ornelasi*, endemic
lagarto alicante de Reid, *Abronia reidi*, endemic
lagarto de cristal cuello simple, *Anguis incomptus*, endemic
lagarto alicante cuello rugoso, *Barisia rudicollis*, endemic
lagartija arenera, lagartija de arena, *Uma exsul*, endemic
lagartija arenera del Colorado, cachora, *Uma notata*, non-endemic
lagartija arenera de Chihuahua, *Uma parphygas*, endemic
huico de Rodeck, *Aspidoscelis rodecki*, endemic
víbora de cascabel, *Crotalus transversus*, endemic
lagartija escofina de Mapimí, *Xantusia bolsonae*, endemic
lagartija nocturna de Sánchez, *Xantusia sanchezi*, endemic
tortuga marina caguama, *Caretta caretta*, non-endemic
tortuga marina verde del Pacífico, tortuga prieta, *Chelonia agassizi*, non-endemic
tortuga marina verde del Atlántico, tortuga blanca, *Chelonia mydas*, non-endemic
tortuga marina de carey, *Eretmochelys imbricata*, non-endemic
tortuga marina escamosa del Atlántico, tortuga lora, *Lepidochelys kempii*, non-endemic
tortuga golfina, tortuga marina escamosa del Pacífico, *Lepidochelys olivacea*, non-endemic
tortuga riverina centroamericana, tortuga blanca, *Dermatemys mawii*, non-endemic
tortuga marina laud, Dermochelys coriácea, non-endemic
tortuga pecho quebrado de Sonoyta, *Kinosternon sonoriense longifemorale*, endemic
tortuga almizclera chopontil, *Claudius angustatus*, non-endemic

galápago de Mapimí, *Gopherus flavomarginatus*, endemic
tortuga de concha blanda negra de Cuatro Ciénegas, *Apalone spinifera atra*, endemic

BIRDS

pato real, *Cairina moschata*, non-endemic
cisne de tundra, *Cygnus columbianus*, non-endemic
colibrí cola hendida, *Doricha eliza*, endemic
colibrí oaxaqueño, *Eupherusa cyanophrys*, endemic
coqueta cresta corta, *Lophornis brachylophus*, endemic
cónedor californiano, *Gymnogyps californianus*, non-endemic
zopilote rey, *Sarcoramphus papa*, non-endemic
cigüeña jabirú, *Jabiru mycteria*, non-endemic
tórtola pecho morado, *Claravis mondetoura*, non-endemic
paloma perdiz tuxtleña, *Geotrygon carrikeri*, endemic
monoto garganta azul, *Aspatha gularis*, non-endemic
momoto pico quilla, *Electron carinatum*, non-endemic
alcuela oscura austral, *Ptychoramphus aleuticus australis*, endemic
mérgado de Craveri, *Synthliboramphus craveri*, non-endemic
ostrero americano, *Haematopus palliatus frazari*, non endemic
chorlo chiflador, *Charadrius melanotos*, non-endemic
charrán embridado guerrerense, *Sterna anaethetus nelsoni*, non-endemic
playero canuto, *Calidris canutus roselaari*, non-endemic
zarapito boreal, *Numenius borealis*, non-endemic
águila cabeza blanca, *Haliaeetus leucocephalus*, non-endemic
águila arpía, *Harpia harpyja*, non-endemic
águila solitaria, *Harpyhaliaetus solitarius*, non-endemic
águila crestada, *Morphnus guianensis*, non-endemic
águila elegante, *Spizaetus ornatus*, non-endemic
águila tirana, *Spizaetus tyrannus*, non-endemic
águila blanquinegra, *Spizastur melanoleucus*, non-endemic
halcón pecho rufo, *Falco deiroleucus*, non-endemic
hocofaisán, *Crax rubra griscomi*, endemic
pavón, guan cornudo, *Oreophasis derbianus*, non-endemic
pajúil, *Penelopina nigra*, non-endemic
codorniz Cotuí, *Colinus virginianus ridgwayi*, endemic
codorniz coluda veracruzana, *Dendrocygna barbata*, endemic
ave sol, *Eurypyga helias*, non-endemic
grulla blanca, *Grus americana*, non-endemic
polluela amarilla, *Coturnicops noveboracensis goldmani*, endemic
polluela negra, *Laterallus jamaicensis coturniculus*, non-endemic
rascón real, *Rallus elegans tenuirostris*, endemic
rascón picudo de Banco Chinchorro, *Rallus longirostris grossi*, endemic
rascón picudo californiano, *Rallus longirostris levipes*, endemic
rascón picudo yucateco, *Rallus longirostris pallidus*
chara azul, *Cyanocorax beecheii*, endemic
chara pinta, *Cyanocorax dickeyi*, endemic
chara garganta blanca, *Cyanolyca mirabilis*, endemic
chara enana, *Cyanolyca nana*, endemic
cascanueces, *Nucifraga columbiana*, non-endemic
trepatroncos gigante de Omiltemi, *Xiphocolaptes promeropirhynchus omiltemensis*, endemic
zacatonero istmeño, *Aimophila sumichrasti*, endemic
semillero azul gris, *Amaurospiza concolor*, non-endemic
juncos ojo oscuro, *Junco hyemalis insularis*, endemic
gorrón cantor de Coronados, *Melospiza melodia coronatorum*, endemic
gorrón indefinido altiplanero or gorrión de Worthen, *Spizella wortheni*, endemic

gorrión serrano, *Xenospiza baileyi*, endemic
pinzón de Guadalupe, *Carpodacus mexicanus amplus*, endemic
pinzón de San Clemente, *Carpodacus mexicanus clementis*, endemic
centzontle de Socorro, *Mimus graysoni*, endemic
cuitlacoche de Cozumel, *Toxostoma guttatum*, endemic
chipe cachete amarillo, *Dendroica chrysoparia*, non-endemic
chipe rosado, *Ergaticus versicolor*, non-endemic
mascarita peninsular, *Geothlypis beldingi*, endemic
mascarita de Altamira, *Geothlypis flavovelata*, endemic
mascarita transvolcánica, *Geothlypis speciosa*, endemic
parula de las Islas Marías, *Parula pitiaayumi insularis*, endemic
reyezuelo de rojo de Guadalupe, *Regulus calendula obscurus*, endemic
tángara chiapaneca, *Tangara cabanisi*, non-endemic
matraca yucateca, *Campylorhynchus yucatanicus*, endemic
chivirín de Nava, *Hylocharis navai*, endemic
chivirín saltarroca de Guadalupe, *Salpinctes obsoletus guadeloupensis*, endemic
mosquero real, *Onychorhynchus coronatus*, non-endemic
vireo gorra negra, *Vireo atricapilla*, non-endemic
vireo de Bell californiano, *Vireo bellii pusillus*, non-endemic
paíño cenizo, *Oceanodroma homochroa*, non-endemic
paíño de Leach de Socorro, *Oceanodroma leucorhoa socorroensis*, non-endemic
paíño de Leach de Coronados, *Oceanodroma leucorhoa willetti*, non-endemic
petrel de Cook, *Pterodroma cookii*
pardela de Revillagigedo, *Puffinus auricularis auricularis*, endemic
pardela mexicana, *Puffinus opisthomelas*, non-endemic
loro nuca amarilla, *Amazona auropalliata*, non-endemic
loro corona azul, *Amazona farinosa*, non-endemic
loro corona lila, *Amazona finschi*, endemic
loro cabeza amarilla, *Amazona oratrix*, non-endemic
loro tamaulipeco, *Amazona viridigenalis*, endemic
guacamaya roja, *Ara macao*, non-endemic
guacamaya verde, *Ara militaris*, non-endemic
periquito de socorro, *Aratinga holochlora brevipes*, endemic
perico del norte, *Aratinga holochlora brewsteri*, endemic
perico catarina de las Islas Marías, *Forpus cyanopygius insularis*, endemic
loro cabeza oscura, *Pionopsitta haematotis*, non-endemic
cotorra serrana occidental, *Rhynchopsitta pachyrhyncha*, endemic
cotorra serrana oriental, *Rhynchopsitta terrisi*, endemic
tecolote canelo, *Aegolius ridgwayi*, non-endemic
tecolote tamaulipeco, *Glaucidium sanchezi*, endemic
tecolote barbudo, *Megascops barbarus*, non-endemic
búho leonado, *Strix fulvescens*, non-endemic
quetzal mesoamericano, *Pharomachrus mocinno*, non-endemic

AMPHIBIANS

rana de Moore, *Lithobates johni*, endemic
rana guerrerense, *Lithobates omiltemanus*, endemic
rana poblana, *Lithobates pueblae*, endemic
rana de Tláloc, *Lithobates tlaloci*, endemic
salamandra ajolote, *Ambystoma mexicanum*, endemic
salamandra saltarina negra, *Ixalotriton niger*, endemic
tritón manchas negras, *Notophthalmus meridionalis*, non-endemic

MAMMALS

Berrendo, *Antilocapra americana*, non-endemic
bura de isla cedros, venado bura, *Odocoileus hemionus cerrosensis*, endemic
bisonte americano, *Bos bison bison*, non-endemic
pecarí de labios blancos, *Tayassu pecari ringens*, non-endemic
ocelote, tigrillo, *Leopardus pardalis*, non-endemic
ocelote, margay, *Leopardus wiedii*, non-endemic
jaguar, tigre, *Panthera onca*, non-endemic
tayra, *Eira barbara*, non-endemic
nutria marina, *Enhydra lutris nereis*, non-endemic
mapache de Islas Marías, *Procyon insulares*, endemic
mapache de Cozumel, *Procyon pygmaeus*, endemic
oso negro, *Ursus americanus eremicus*, non-endemic
ballena franca, *Eubalaena japonica*, non-endemic
vaquita, *Phocoena sinus*, endemic
armadillo centroamericano, armadillo rabo liso norteño, *Cabassous centrales*, non-endemic
murciélagos platanero, *Musonycteris harrisoni*, endemic
vampiro falso de Linneo, *Vampyrum spectrum*, non-endemic
miotis cabeza plana, *Myotis planiceps*, endemic
miotis pescador, *Myotis vivesi*, endemic
tlacuache de agua, *Chironectes minimus*, non-endemic
topo occidental, *Scalopus aquaticus*, non-endemic
topo pata ancha, *Scapanus latimanus anthonyi*, non-endemic
teporingo, conejo de los volcanes, *Romerolagus diazi*, endemic
conejo matorralero de la Isla Cedros, *Sylvilagus bachmani cerrosensis*, endemic
conejo de Tres Marías, *Sylvilagus graysoni*, endemic
conejo de Omiltemi, *Sylvilagus insonus*, endemic
conejo de San José, *Sylvilagus mansuetus*, endemic
tapir centroamericano, *Tapirus bairdii*, non-endemic
oso hormiguero dorado, *Cyclopes didactylus*, non-endemic
oso hormiguero, brazo fuerte, tamandúa norteño, *Tamandua mexicana hesperia*, endemic
oso hormiguero, brazo fuerte, tamandúa norteño, *Tamandua mexicana mexicana*, non-endemic
lobo fino de Guadalupe, *Arctocephalus townsendi*, endemic
mono aullador, saraguato de manto, *Alouatta palliata*, non-endemic
mono aullador, saraguato yucateco, *Alouatta pigra*, non-endemic
mono araña, *Ateles geoffroyi*, non-endemic
castor, *Castor canadensis*, non-endemic
puerco espín del norte, *Erethizon dorsatum*, non-endemic
tuza michoacana, *Zygogeomys trichopus*, endemic
ratón de abazones sonorense de Monserrat, *Chaetodipus baileyi insulares*, endemic
ratón de abazones de San José, *Chaetodipus spinatus bryanti*, endemic
ratón de abazones de San Francisco, *Chaetodipus spinatus latijugularis*, endemic
rata canguro de San José, *Dipodomys insulares*, endemic
rata canguro de Margarita, *Dipodomys margaritae*, endemic
meteoro de California, *Microtus californicus*, non-endemic
meteoro de prado, *Microtus pennsylvanicus*, non-endemic
rata cambalachera de San Martín, *Neotoma martinensis*, endemic
rata cambalachera de Turner, *Neotoma varia*, endemic
ratón de Ángel de la Guarda, *Peromyscus guardia*, endemic
perrito de las praderas, perro llanero mexicano, *Cynomys mexicanus*, endemic
manatí del Caribe, *Trichechus manatus*, non-endemic
musaraña de Arizona, *Sorex arizonae*, non-endemic